

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 September 2005 (29.09.2005)

PCT

(10) International Publication Number
WO 2005/089118 A2

(51) International Patent Classification: Not classified

PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA,
ZM, ZW.

(21) International Application Number:
PCT/US2005/006925

(22) International Filing Date: 3 March 2005 (03.03.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/550,154 4 March 2004 (04.03.2004) US

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE,
EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS,
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM,
PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY,
TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA,
ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ,
NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM,
AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT,
BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,
HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG)

— as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(71) Applicant (for all designated States except US): BRIS-
TOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box
4000, Route 206 and ProvinceLine Road, Princeton, New
Jersey 08543-4000 (US).


(72) Inventors; and

(75) Inventors/Applicants (for US only): LI, James J.
[US/US]; 76 Stanford Road East, Pennington, NJ 08534
(US). HAMANN, Lawrence [US/US]; 24 East Riding
Drive, Cherry Hill, NJ 08003 (US). AUGERI, David
[US/US]; 107 Carter Road, Princeton, NJ 08540 (US).
BI, Yingzhi [US/US]; 5 Mershon Lane, Plainsboro, New
Jersey 08536 (US).

(74) Agents: DUNCAN, Sammy G. et al.; Bristol-Myers
Squibb Company, P.O. Box 4000, Princeton, New Jersey
08543-4000 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,

(54) Title: NOVEL BICYCLIC COMPOUNDS AS MODULATORS OF ANDROGEN RECEPTOR FUNCTION AND METHOD



(I)

(57) Abstract: The present invention relates to bicyclic compounds according to formula
(I), pharmaceutical compositions containing such compounds and methods of using such
compounds in the treatment of androgen receptor-associated conditions, such as age-related
diseases, for example sarcopenia, formula (I) wherein R1, R2, R2' R5, G, n and W are de-
fined herein.

WO 2005/089118 A2

NOVEL BICYCLIC COMPOUNDS AS MODULATORS OF ANDROGEN RECEPTOR FUNCTION AND METHOD

RELATED APPLICATIONS

- 5 [0001] This application claims priority benefit under Title 35 § 119(e) of United States provisional Application No. 60/550,154, filed March 4, 2004, the contents of which are herein incorporated by reference.

FIELD OF THE INVENTION

- 10 [0002] The present invention relates to bicyclic compounds, pharmaceutical compositions containing such compounds and methods of using such compounds in the treatment of androgen receptor-associated conditions, such as age-related diseases, for example sarcopenia.

BACKGROUND OF THE INVENTION

- 15 [0003] Nuclear hormone receptors (NHR's) constitute a large super-family of structurally-related and sequence-specific gene regulators scientists have named "ligand-dependent transcription factors." R.M. Evans, *Science*, 240:889 (1988). The steroid binding NHR's (SB-NHR's) form a recognized subset of the NHR's, including
20 the progesterone receptor (PR), androgen receptor (AR), estrogen receptor (ER), glucocorticoid receptor (GR) and mineralocorticoid receptor (MR). The conventional nuclear hormone receptors are generally transactivators in the presence of ligand, which selectively bind to the NHR in a way that effects gene transcription. In the absence of a corresponding ligand, some of the orphan receptors behave as if they are
25 transcriptionally inert. Others, however, behave as either constitutive activators or repressors. These orphan nuclear hormone receptors are either under the control of ubiquitous ligands that have not been identified, or do not need to bind ligand to exert these activities.

- [0004] The AR is a ligand-activated transcriptional regulatory protein that
30 mediates induction of male sexual development and function through its activity with endogenous androgens. In addition, androgens are associated with male and female maintenance of muscle mass and strength, bone mass and erythropoiesis. Androgens,

such as testosterone, also play an important role in many physiological processes, such as differentiation of male internal and external genitalia, development and maintenance of male secondary sexual characteristics (e.g., the development of prostate, seminal vesicles, penis, scrotum, skeletal muscle, redistribution of body fat, stimulation of long bone growth, closure of epiphyses, development of male hair growth pattern and enlargement of larynx), the maintenance of sexual behavior and function (e.g., libido and potency) and spermatogenesis (in man).

[0005] As one ages, the serum androgen concentration in the body declines. The age dependent decline in androgens is associated with changes in body composition for men and women, such as a lower percentage of muscle mass and an increase in body fat, e.g., sarcopenia. In this regard, modulation of the AR gene can have an impact on the physiological effects associated with androgen production. However, the effectiveness of known modulators of steroid receptors is often tempered by their undesired side-effect profile, particularly during long-term administration. For example, the administration of synthetic androgens has been associated with liver damage, prostate cancer, adverse effects on male sexual function and adverse effects associated with cardiovascular and erythropoietic function.

[0006] Numerous synthetically-derived steroidal and non-steroidal agonists and antagonists have been described for the members of the SB-NHR family. Many of these agonist and antagonist ligands are used clinically in man to treat a variety of medical conditions. RU486 (mifepristone) is an example of a synthetic antagonist of the PR, which is utilized as a birth control agent (Vegeto et al., *Cell* **69**: 703-713 (1992)). Flutamide is an example of an antagonist of the AR, which is utilized for the treatment of prostate cancer (Neri et al, *Endo.* **91**, 427-437 (1972)). Tamoxifen is an example of a tissue-selective modulator of the ER function, that is used in the treatment of breast cancer (Smigel *J. Natl. Cancer Inst.* **90**, 647-648 (1998)). Tamoxifen can function as an antagonist of the ER in breast tissue while acting as an agonist of the ER in bone (Grese et al., *Proc. Natl. Acad. Sci. USA* **94**, 14105-14110 (1997)). Because of the tissue-selective effects seen for Tamoxifen, this agent, and agents like it, are referred to as tissue-selective estrogen receptor modulators. In addition to synthetically-derived non-endogenous ligands, non-endogenous ligands for NHR's can be obtained from food sources (Regal et al., *Proc. Soc. Exp. Biol. Med.*

223, 372-378 (2000) and Hempstock et al., *J. Med. Food* 2, 267-269 (1999)). The flavanoid phytoestrogens are an example of an unnatural ligand for SB-NHR's that are readily obtained from a food source such as soy (Quella et al., *J. Clin. Oncol.* 18, 1068-1074 (2000) and Banz et al., *J. Med. Food* 2, 271-273 (1999)). The ability to
5 modulate the transcriptional activity of an individual NHR by the addition of a small molecule ligand, makes these receptors ideal targets for the development of pharmaceutical agents for a variety of disease states.

[0007] As mentioned above, non-natural ligands can be synthetically engineered to serve as modulators of the function of NHR's. In the case of SB-NHR's,
10 engineering of an unnatural ligand can include the identification of a core structure which mimics the natural steroid core system. This can be achieved by random screening against several SB-NHR's, or through directed approaches using the available crystal structures of a variety of NHR ligand binding domains (Bourguet et al., *Nature* 375, 377-382 (1995), Brzozowski, et al., *Nature* 389, 753-758 (1997),
15 Shiau et al., *Cell* 95, 927-937 (1998) and Tanenbaum et al., *Proc. Natl. Acad. Sci. USA* 95, 5998-6003 (1998)). Differential substitution about such a steroid mimic core can provide agents with selectivity for one receptor versus another. In addition, such modifications can be employed to obtain agents with agonist or antagonist activity for a particular SB-NHR. Differential substitution about the steroid mimic core can result
20 in the formation of a series of high affinity agonists and antagonists with specificity for, for example, ER versus PR versus AR versus GR versus MR. Such an approach of differential substitution has been reported, for example, for quinoline based modulators of steroid NHRs in Hamann et. al., *J. Med. Chem.*, 41, 623 (1998); Hamann et. al., *J. Med. Chem.* 42, 210 (1999); WO 9749709; US 5696133; US
25 5696130; US 5696127; US 5693647; US 5693646; US 5688810; US 5688808 and WO 9619458, all incorporated herein by reference.


[0008] Accordingly, identification of compounds which have good specificity for one or more steroid receptors, but which have reduced or no cross-reactivity for other steroid or intracellular receptors, would be of significant value in the treatment of
30 male and female hormone-responsive diseases. There is, therefore, a need in the art for the identification of selective modulators of the steroid binding nuclear hormone receptors, particularly non-steroidal, non-toxic tissue selective androgen receptor

modulators, which activate the androgen receptor in skeletal muscle while demonstrating limited or neutral effect on other androgen responsive (e.g., prostate) tissues.

5

SUMMARY OF THE INVENTION

[0009] In a first embodiment of the present invention, compounds are provided which are capable of modulating the function of a nuclear hormone receptor having the general formula I



10

I

wherein

R_1 is independently selected from the group consisting of H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, arylalkyl, substituted arylalkyl, CO_2R_4 , CONR_4R_4 , and CH_2OR_4 ;

R_2 is independently selected from the group consisting of H, alkyl, substituted alkyl, OR_3 , SR_3 , halo, NHR_3 , NHCOR_4 , NHCO_2R_4 , $\text{NHCONR}_4\text{R}_4$ and NHSO_2R_4 ;

R_2' is independently selected from the group consisting of OR_3 , SR_3 , halo, NHR_3 , NHCOR_4 , NHCO_2R_4 , $\text{NHCONR}_4\text{R}_4$ and NHSO_2R_4 ;

R_3 is independently selected from the group consisting of H, alkyl, substituted alkyl, CHF_2 , CF_3 and COR_4 ;

R_4 , is independently selected from the group consisting of H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

R_5 is independently selected from the group consisting of H, OR_3 , CF_3 , alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl

and substituted heteroaryl, wherein R_5 and R_5 taken together can form a double bond with O, S, NR_7 or CR_7R_7 when \parallel is a single bond, it being understood that only one R_5 is present when \parallel is a double bond;

W is CR_6R_6 , SO, or SO_2 ;

5 R_6 is independently selected from the group consisting of H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl, wherein at least one R_6 is H, or R_6 and R_6 taken together can form a double bond with O, S, NR_7 or CR_7R_7 ;

10 R_7 is independently selected from the group consisting of H, OR_4 , alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

G is selected from the group consisting of aryl, heterocyclo and heteroaryl,
15 wherein said aryl, heterocyclo or heteroaryl is mono- or polycyclic, and is optionally substituted with one or more substituents selected from the group consisting of hydrogen, halo, CN, CF_3 , OR_4 , CO_2R_4 , NR_4R_4 , $CONR_4R_4$, CH_2OR_4 , alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and
20 substituted heteroaryl; and

n is 1 or 2.

[0010] The definition of formula I above includes all prodrug esters, stereoisomers and pharmaceutically acceptable salts of formula I.

25

[0011] A preferred embodiment of the present invention provides compounds of formula I wherein

R_1 is H or alkyl;

R_2' is OH;


30 R_5 is alkyl or substituted alkyl;

\parallel is a double bond;

W is C=O; and

n is 1.

- 5 [0012] Further preferred embodiments include compounds of formula I where G is selected from:



- 10 wherein

R_8 , R_9 , R_{10} and R_{11} are independently selected from the group consisting of H, NO_2 , CN, CF_3 , OR_4 , CO_2R_4 , NR_4R_4 , CONR_4R_4 , CH_2OR_4 , alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and

- 15 substituted heteroaryl;

A, B, C, D, E and F are independently selected from the group consisting of N and CR_{12} ;

J, K, L, P and Q are independently selected from the group consisting of NR_{13} , O, S, SO, SO_2 and $\text{CR}_{13}\text{R}_{13}$;

- 20 R_{12} is independently selected from the group consisting of a direct bond and R_{13} ;

R_{13} is independently selected from the group consisting of H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, arylalkyl, substituted arylalkyl, CO_2R_4 , CONR_4R_4 and CH_2OR_4 ; and

- 25 z is 0 or 1.

[0013] A further preferred embodiment of the present invention provides compounds wherein R₈ is CN.

[0014] The compounds of formula I modulate the function of the nuclear hormone
5 receptors, particularly the androgen receptor, and include compounds which are, for
example, selective agonists, partial agonists, antagonists or partial antagonists of the
androgen receptor. Preferably the compounds of formula I possess activity as agonists
of the androgen receptor and may be used in the treatment of diseases or disorders
associated with androgen receptor activity, such as maintenance of muscle strength
10 and function (e.g., in the elderly); reversal or prevention of frailty or age-related
functional decline ("ARFD") in the elderly (e.g., sarcopenia); prevention of catabolic
side effects of glucocorticoids; prevention and treatment of reduced bone density or
growth (e.g., osteoporosis and osteopenia); treatment of chronic fatigue syndrome
(CFS); chronic myalgia; treatment of acute fatigue syndrome and muscle loss
15 following elective surgery (e.g., post-surgical rehabilitation); treatment of urinary
incontinence, acceleration of wound healing; accelerating bone fracture repair (such as
accelerating the recovery of hip fracture patients); treatment of wasting secondary to
fractures and wasting in connection with chronic obstructive pulmonary disease
(COPD), chronic liver disease, AIDS, weightlessness, cancer cachexia, burn and
20 trauma recovery, chronic catabolic state (e.g., coma), eating disorders (e.g., anorexia)
and chemotherapy.

[0015] In a second embodiment of the present invention, pharmaceutical
compositions, comprising at least one compound according to formula I and at least
one pharmaceutically acceptable diluent or carrier are provided.

25 [0016] In a third embodiment of the present invention, methods for preventing,
inhibiting or treating the progression or onset of diseases or disorders associated with
nuclear hormone receptors, comprising administering to a patient in need a
therapeutically effective amount of a compound according to formula I are provided.

[0017] In a preferred embodiment of the present invention, methods are provided,
30 wherein the nuclear hormone receptor is the androgen receptor.

[0018] In a further preferred embodiment of the present invention, methods are
provided wherein the disease or disorder associated with the androgen receptor is

selected from the group consisting of maintenance of muscle strength and function; reversal or prevention of frailty or age-related functional decline ("ARFD") in the elderly (e.g., sarcopenia); prevention of catabolic side effects of glucocorticoids; prevention and treatment of reduced bone density or growth (e.g., osteoporosis and osteopenia); treatment of chronic fatigue syndrome (CFS); chronic myalgia; treatment of acute fatigue syndrome and muscle loss following elective surgery (e.g., post-surgical rehabilitation); treatment of urinary incontinence, acceleration of wound healing; accelerating bone fracture repair (such as accelerating the recovery of hip fracture patients); treatment of wasting secondary to fractures and wasting in connection with chronic obstructive pulmonary disease (COPD), chronic liver disease, AIDS, weightlessness, cancer cachexia, burn and trauma recovery, chronic catabolic state (e.g., coma), eating disorders (e.g., anorexia) and chemotherapy.

[0019] In a still further preferred embodiment of the present invention, methods further comprising a therapeutically effective amount of at least one other therapeutic agent are provided.

DETAILED DESCRIPTION OF THE INVENTION

Abbreviations

[0020] The following abbreviations are employed herein:

Chiralpak® = Trademark of Chiral Technologies, Inc. Eaton, PA

DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene

AcOH = acetic acid

EtOAc = ethyl acetate

HPLC = high performance liquid chromatography

MeOH = methanol

MS or Mass Spec = mass spectrometry

YMC® = trademark of YMC Co, Ltd., Kyoto, Japan

CuBr = copper(I) bromide

CuI = copper(I) iodide

CuCN = copper(I) cyanide

CsF = cesium fluoride

Et₃N = triethylamine

LDA = lithium diisopropylamide

NMP = 1-methyl-2-pyrrolidinone

KOH = potassium hydroxide

Pd/C = palladium on activated charcoal

5 PyBrop = bromo-trispyrrolidino-phosphonium hexafluorophosphate

TFA = trifluoroacetic acid

THF = tetrahydrofuran

mp. = melting point

min = minute(s)

10 h = hour(s)

L = liter

mL = milliliter

μ L = microliter

g = gram(s)

15 mg = milligram(s)

mol = moles

mmol = millimole(s)

nM = nanomolar

rt or RT = room temperature

20

Definitions

[0021] The following definitions apply to the terms as used throughout this specification, unless otherwise limited in specific instances.

[0022] As used herein, the term "alkyl" denotes branched or unbranched
25 hydrocarbon chains, preferably having about 1 to about 8 carbons, such as, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, 2-methylpentyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethyl pentyl, octyl, 2,2,4-trimethylpentyl and the like. "Substituted alkyl" includes an alkyl group optionally substituted with one or more functional groups which are attached commonly to such chains, such as,
30 hydroxyl, bromo, fluoro, chloro, iodo, mercapto or thio, cyano, alkylthio, heterocyclyl, aryl, heteroaryl, carboxyl, carbalkoyl, alkyl, alkenyl, nitro, amino, alkoxyl, amido, and

the like to form alkyl groups such as trifluoro methyl, 3-hydroxyhexyl, 2-carboxypropyl, 2-fluoroethyl, carboxymethyl, cyanobutyl and the like.

[0023] Unless otherwise indicated, the term "cycloalkyl" as employed herein alone or as part of another group includes saturated or partially unsaturated (containing 1 or more double bonds) cyclic hydrocarbon groups containing 1 to 3 rings, including monocyclicalkyl, bicyclicalkyl and tricyclicalkyl, containing a total of 3 to 20 carbons forming the rings, preferably 3 to 10 carbons, forming the ring and which may be fused to 1 or 2 aromatic rings as described for aryl, which include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl and cyclododecyl, cyclohexenyl,



[0024] "Substituted cycloalkyl" includes a cycloalkyl group optionally substituted with 1 or more substituents such as halogen, alkyl, alkoxy, hydroxy, aryl, aryloxy, arylalkyl, cycloalkyl, alkylamido, alkanoylamino, oxo, acyl, arylcarbonylamino, amino, nitro, cyano, thiol and/or alkylthio and/or any of the substituents included in the definition of "substituted alkyl."


[0025] Unless otherwise indicated, the term "alkenyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons, and more preferably 2 to 8 carbons in the normal chain, which include one or more double bonds in the normal chain, such as vinyl, 2-propenyl, 3-butenyl, 2-butenyl, 4-pentenyl, 3-pentenyl, 2-hexenyl, 3-hexenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 3-octenyl, 3-nonenyl, 4-decenyl, 3-undecenyl, 4-dodecenyl, 4,8,12-tetradecatrienyl, and the like. "Substituted alkenyl" includes an alkenyl group optionally substituted with one or more substituents, such as the substituents included above in the definition of "substituted alkyl" and "substituted cycloalkyl."


[0026] Unless otherwise indicated, the term "alkynyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons and more preferably 2 to 8 carbons in the normal chain, which include one or more triple bonds in the normal chain, such as 2-propynyl, 3-butynyl, 2-butynyl, 4-pentynyl, 3-pentynyl, 2-hexynyl, 3-hexynyl, 2-heptynyl, 3-heptynyl, 4-heptynyl, 3-octynyl, 3-nonyl, 4-decynyl, 3-undecynyl, 4-dodecynyl and the like. "Substituted alkynyl" includes an alkynyl group optionally substituted with one or more substituents, such as the substituents included above in the definition of "substituted alkyl" and "substituted cycloalkyl."

[0027] The terms "arylalkyl", "arylalkenyl" and "arylalkynyl" as used alone or as part of another group refer to alkyl, alkenyl and alkynyl groups as described above having an aryl substituent. Representative examples of arylalkyl include, but are not limited to, benzyl, 2-phenylethyl, 3-phenylpropyl, phenethyl, benzhydryl and naphthylmethyl and the like. "Substituted arylalkyl" includes arylalkyl groups wherein the aryl portion is optionally substituted with one or more substituents, such as the substituents included above in the definition of "substituted alkyl" and "substituted cycloalkyl."

[0028] The term "halogen" or "halo" as used herein alone or as part of another group refers to chlorine, bromine, fluorine, and iodine.


[0029] Unless otherwise indicated, the term "aryl" or "Ar" as employed herein alone or as part of another group refers to monocyclic and polycyclic aromatic groups containing 6 to 10 carbons in the ring portion (such as phenyl or naphthyl including 1-naphthyl and 2-naphthyl) and may optionally include one to three additional rings fused to a carbocyclic ring or a heterocyclic ring (such as aryl, cycloalkyl, heteroaryl or cycloheteroalkyl rings), for example






[0030] "Substituted aryl" includes an aryl group optionally substituted with one or more functional groups, such as halo, haloalkyl, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, trifluoromethyl, trifluoromethoxy, alkynyl, cycloalkyl-alkyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, aryloxy, aryloxyalkyl, arylalkoxy, alkoxycarbonyl, arylcarbonyl, arylalkenyl, aminocarbonylaryl, arylthio, arylsulfinyl, arylazo, heteroarylalkyl, heteroarylalkenyl, heteroarylheteroaryl, heteroaryloxy, hydroxy, nitro, cyano, amino, substituted amino wherein the amino includes 1 or 2 substituents (which are alkyl, aryl or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkoxyarylthio, alkylcarbonyl, arylcarbonyl, alkylaminocarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonylamino or arylsulfonaminocarbonyl and/or any of the alkyl substituents set out herein.

[0031] Unless otherwise indicated, the term "heteroaryl" as used herein alone or as part of another group refers to a 5- or 7- membered aromatic ring which includes 1, 2, 3 or 4 hetero atoms such as nitrogen, oxygen or sulfur and such rings fused to an aryl, cycloalkyl, heteroaryl or heterocycloalkyl ring (e.g. benzothiophenyl, indolyl), and includes possible N-oxides. "Substituted heteroaryl" includes a heteroaryl group optionally substituted with 1 to 4 substituents, such as the substituents included above in the definition of "substituted alkyl" and "substituted cycloalkyl." Examples of heteroaryl groups include the following:





5

and the like.

[0032] The term “heterocyclo”, heterocycle or heterocyclic ring, as used herein, represents an unsubstituted or substituted stable 5- to 7-membered monocyclic ring system which may be saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from N, O or S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyclic groups include, but is not limited to, piperidinyl, piperazinyl, oxopiperazinyl, oxopiperidinyl, oxopyrrolidinyl, oxoazepinyl, azepinyl, pyrrolyl, pyrrolidinyl, furanyl, thienyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazolyl, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isooxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, thiadiazolyl, tetrahydropyranyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, and oxadiazolyl.

[0033] The compounds of formula I can be present as salts, which are also within the scope of this invention. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred. If the compounds of formula I have, for example, at least one basic center, they can form acid addition salts. These are formed, for example, with strong inorganic acids, such as mineral acids, for example sulfuric acid, phosphoric acid or a hydrohalic acid, with strong organic carboxylic

acids, such as alkanecarboxylic acids of 1 to 4 carbon atoms which are unsubstituted or substituted, for example, by halogen, for example acetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, (for example aspartic or glutamic acid or lysine or arginine), or benzoic acid, or with organic sulfonic acids, such as (C₁-C₄) alkyl or arylsulfonic acids which are unsubstituted or substituted, for example by halogen, for example methyl- or *p*-toluene- sulfonic acid. Corresponding acid addition salts can also be formed having, if desired, an additionally present basic center. The compounds of formula I having at least one acid group (for example COOH) can also form salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine, piperidine, pyrrolidine, a mono, di or tri-lower alkylamine, for example ethyl, *tert*-butyl, diethyl, diisopropyl, triethyl, tributyl or dimethyl-propylamine, or a mono, di or trihydroxy lower alkylamine, for example mono, di or triethanolamine. Corresponding internal salts may furthermore be formed. Salts which are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free compounds of formula I or their pharmaceutically acceptable salts, are also included.

[0034] Preferred salts of the compounds of formula I which contain a basic group include monohydrochloride, hydrogensulfate, methanesulfonate, phosphate or nitrate.

[0035] Preferred salts of the compounds of formula I which contain an acid group include sodium, potassium and magnesium salts and pharmaceutically acceptable organic amines.

[0036] The term "modulator" refers to a chemical compound with capacity to either enhance (e.g., "agonist" activity) or inhibit (e.g., "antagonist" activity) a functional property of biological activity or process (e.g., enzyme activity or receptor binding); such enhancement or inhibition may be contingent on the occurrence of a specific event, such as activation of a signal transduction pathway, and/or may be manifest only in particular cell types.

[0037] The term "prodrug esters" as employed herein includes esters and carbonates formed by reacting one or more hydroxyls of compounds of formula I with alkyl, alkoxy, or aryl substituted acylating agents employing procedures known to those skilled in the art to generate acetates, pivalates, methylcarbonates, benzoates and the like.

[0038] Any compound that can be converted in vivo to provide the bioactive agent (i.e., the compound of formula I) is a prodrug within the scope and spirit of the invention.

[0039] Various forms of prodrugs are well known in the art. A comprehensive description of prodrugs and prodrug derivatives are described in:

(a) *The Practice of Medicinal Chemistry*, Camille G. Vermuth et al., Ch 31, (Academic Press, 1996);

(b) *Design of Prodrugs*, edited by H. Bundgaard, (Elsevier, 1985);

(c) *A Textbook of Drug Design and Development*, P. Krosgaard-Larson and H. Bundgaard, eds., Ch. 5, pgs, 113-191 (Harwood Academic Publishers, 1991). Said references are incorporated herein by reference.


[0040] An administration of a therapeutic agent of the invention includes administration of a therapeutically effective amount of the agent of the invention. The term "therapeutically effective amount" as used herein refers to an amount of a therapeutic agent to treat or prevent a condition treatable by administration of a composition of the invention. That amount is the amount sufficient to exhibit a detectable therapeutic or preventative or ameliorative effect. The effect may include, for example, treatment or prevention of the conditions listed herein. The precise effective amount for a subject will depend upon the subject's size and health, the nature and extent of the condition being treated, recommendations of the treating physician, and the therapeutics or combination of therapeutics selected for administration. Thus, it is not useful to specify an exact effective amount in advance.

[0041] All stereoisomers of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure form. The compounds of the present invention can have asymmetric centers at any of the carbon atoms including any one of the R substituents. Consequently, compounds of formula I can exist in enantiomeric or diastereomeric forms or in mixtures thereof. The

processes for preparation can utilize racemates, enantiomers or diastereomers as starting materials. When diastereomeric or enantiomeric products are prepared, they can be separated by conventional methods for example, chromatographic, chiral HPLC or fractional crystallization.

- 5 [0042] The compounds of formula I of the present invention can be prepared as shown in the following reaction schemes and description thereof, as well as relevant published literature procedures that may be used by one skilled in the art. Exemplary reagents and procedures for these reactions appear hereinafter and in the working Examples.
- 10 [0043] Chemistry used to synthesize preferred compounds of the general structure I is shown below in Schemes I–IV.


SCHEME I



[0044] Scheme I describes methods for preparing compounds of formula Ia and Ib (a subset of compound I) from N-protected amino acids ester of formula II, which can be obtained commercially or can be prepared by methods known in the literature


or can be readily prepared by one skilled in the art. An intermediate of formula **II** is treated with a reducing agent, such as super hydride, to form an alcohol intermediate **III**, which can be oxidized to an aldehyde intermediate **IV**. Subsequently **IV** can be treated with a Grignard reagent or an organolithium reagent to form an alcohol intermediate **V** which can be oxidized to a ketone intermediate **VI**. Removal of N-protecting group (PG) can be achieved by an acid such as TFA in dichloromethane or other methods known in the literature or by one skilled in the art to provide an intermediate of formula **VII**. Amide intermediate **IX** can be obtained by reaction of intermediate **VII** and an acid intermediate **VIII** (Scheme III and IIIa). Cyclization can be carried out with a base such as piperidine to give compound **Ia** which can be dehydrated to give compound **Ib**.

SCHEME II




[0045] Scheme II describes methods to prepare compounds of formula **Ic** and **Id** (a subset of compound **I**) from amine intermediate **VII** (Scheme I). Reaction of intermediate **VII** and sulfonyl chloride **X** (Scheme IV) in the presence of a base such as triethylamine to give sulfonamide intermediate **XI**. Cyclization can be carried out with a base such as piperidine to give compound **Ic** which can be dehydrated to give compound **Id**.

SCHEME III



- 5 [0046] Scheme III describes methods to prepare compounds of formula **VIII** from alcohol **XII**, which can be obtained commercially or can be prepared by methods known in the literature or can be readily prepared by one skilled in the art. Alcohol can be converted to halides **XIII** by methods known in the literature or can react with mesyl chloride in the presence of a base to give mesylate intermediate **XIII**. Reaction of intermediate **XIII** and potassium cyanide followed by hydrolysis give the acid intermediate **VIII**.


SCHEME IIIa



- 15 [0047] Scheme IIIa describes methods to prepare compounds of formula **VIIIa** (a subset of compound **VIII**). Intermediates **XIV** are treated with tert-butylnitrite or an equivalent reagent in the presence of a halide source such as CuI or CuBr to afford an aryl or heteroaryl halide of formula **XV** or **XVI**. Either **XV** or **XVI** can be reacted with diethyl malonate in the presence of sodium hydride and CuI to give intermediate


XVII. Basic hydrolysis followed by de-carboxylation under acidic condition provide compound **VIIIa**.

SCHEME IIIb



[0048] Scheme IIIb describes additional methods to prepare compounds of formula **VIIIa** (a subset of compound **VIII**). Compound **XVIII** can react with NBS to give a bromo-intermediate **XIX**. Reaction of intermediate **XIX** and potassium cyanide followed by hydrolysis give acid intermediate **VIIIa**. Alternatively, Compound **XVIII** can react with a base such LDA followed by ClCO_2Me to give an ester intermediate **XX**. Hydrolysis of ester **XX** under basic condition provide acid intermediate **VIIIa**.

SCHEME IV



[0049] Scheme IV describes methods for preparing sulfonyl chloride **X** from an alkyl chloride precursor **XXI**. An alkyl halide was first treated with sodium thiosulfate, then reacted with chlorine gas to give sulfonyl chloride **X**. Additional

methods for preparing sulfonyl chloride X can be found in the literature or be readily carried out by one skilled in the art.

USE AND UTILITY

5 A. Utilities

[0050] The compounds of the present invention modulate the function of the nuclear hormone receptors, particularly the androgen receptor, and include compounds which are, for example, selective agonists, partial agonists, antagonists or partial antagonists of the androgen receptor (AR). Thus, the present compounds are
10 useful in the treatment of AR-associated conditions. An "AR-associated condition," as used herein, denotes a condition or disorder which can be treated by modulating the function or activity of an AR in a subject, wherein treatment comprises prevention, partial alleviation or cure of the condition or disorder. Modulation may occur locally, for example, within certain tissues of the subject, or more extensively throughout a
15 subject being treated for such a condition or disorder.

[0051] The compounds of the present invention can be administered to animals, preferably humans, for the treatment of a variety of conditions and disorders, including, but not limited to maintenance of muscle strength and function (e.g., in the elderly); reversal or prevention of frailty or age-related functional decline ("ARFD")
20 in the elderly (e.g., sarcopenia); treatment of catabolic side effects of glucocorticoids; prevention and/or treatment of reduced bone mass, density or growth (e.g., osteoporosis and osteopenia); treatment of chronic fatigue syndrome (CFS); chronic myalgia; treatment of acute fatigue syndrome and muscle loss following elective surgery (e.g., post-surgical rehabilitation); accelerating of wound healing; accelerating
25 bone fracture repair (such as accelerating the recovery of hip fracture patients); accelerating healing of complicated fractures, e.g. distraction osteogenesis; in joint replacement; prevention of post-surgical adhesion formation; acceleration of tooth repair or growth; maintenance of sensory function (e.g., hearing, sight, olfaction and taste); treatment of periodontal disease; treatment of wasting secondary to fractures
30 and wasting in connection with chronic obstructive pulmonary disease (COPD), chronic liver disease, AIDS, weightlessness, cancer cachexia, burn and trauma recovery, chronic catabolic state (e.g., coma), eating disorders (e.g., anorexia) and

chemotherapy; treatment of cardiomyopathy; treatment of thrombocytopenia;
treatment of growth retardation in connection with Crohn's disease; treatment of short
bowel syndrome; treatment of irritable bowel syndrome; treatment of inflammatory
bowel disease; treatment of Crohn's disease and ulcerative colitis; treatment of
5 complications associated with transplantation; treatment of physiological short stature
including growth hormone deficient children and short stature associated with chronic
illness; treatment of obesity and growth retardation associated with obesity; treatment
of anorexia (e.g., associated with cachexia or aging); treatment of hypercortisolism
and Cushing's syndrome; Paget's disease; treatment of osteoarthritis; induction of
10 pulsatile growth hormone release; treatment of osteochondrodysplasias; treatment of
depression, nervousness, irritability and stress; treatment of reduced mental energy
and low self-esteem (e.g., motivation/assertiveness); improvement of cognitive
function (e.g., the treatment of dementia, including Alzheimer's disease and short
term memory loss); treatment of catabolism in connection with pulmonary
15 dysfunction and ventilator dependency; treatment of cardiac dysfunction (e.g.,
associated with valvular disease, myocardial infarction, cardiac hypertrophy or
congestive heart failure); lowering blood pressure; protection against ventricular
dysfunction or prevention of reperfusion events; treatment of adults in chronic
dialysis; reversal or slowing of the catabolic state of aging; attenuation or reversal of
20 protein catabolic responses following trauma (e.g., reversal of the catabolic state
associated with surgery, congestive heart failure, cardiac myopathy, burns, cancer,
COPD etc.); reducing cachexia and protein loss due to chronic illness such as cancer
or AIDS; treatment of hyperinsulinemia including nesidioblastosis; treatment of
immunosuppressed patients; treatment of wasting in connection with multiple
25 sclerosis or other neurodegenerative disorders; promotion of myelin repair;
maintenance of skin thickness; treatment of metabolic homeostasis and renal
homeostasis (e.g., in the frail elderly); stimulation of osteoblasts, bone remodeling and
cartilage growth; regulation of food intake; treatment of insulin resistance, including
NIDDM, in mammals (e.g., humans); treatment of insulin resistance in the heart;
30 improvement of sleep quality and correction of the relative hyposomatotropism of
senescence due to high increase in REM sleep and a decrease in REM latency;
treatment of hypothermia; treatment of congestive heart failure; treatment of

lipodystrophy (e.g., in patients taking HIV or AIDS therapies such as protease inhibitors); treatment of muscular atrophy (e.g., due to physical inactivity, bed rest or reduced weight-bearing conditions); treatment of musculoskeletal impairment (e.g., in the elderly); improvement of the overall pulmonary function; treatment of sleep disorders; and the treatment of the catabolic state of prolonged critical illness; treatment of hirsutism, acne, seborrhea, androgenic alopecia, anemia, hyperpilosity, benign prostate hypertrophy, adenomas and neoplasies of the prostate (e.g., advanced metastatic prostate cancer) and malignant tumor cells containing the androgen receptor, such as is the case for breast, brain, skin, ovarian, bladder, lymphatic, liver and kidney cancers; cancers of the skin, pancreas, endometrium, lung and colon; osteosarcoma; hypercalcemia of malignancy; metastatic bone disease; treatment of spermatogenesis, endometriosis and polycystic ovary syndrome; counteracting preeclampsia, eclampsia of pregnancy and preterm labor; treatment of premenstrual syndrome; treatment of vaginal dryness; age related decreased testosterone levels in men, male menopause, hypogonadism, male hormone replacement, male and female sexual dysfunction (e.g., erectile dysfunction, decreased sex drive, sexual well-being, decreased libido), urinary incontinence, male and female contraception, hair loss, Reaven's Syndrome and the enhancement of bone and muscle performance/strength. The term treatment is also intended to include prophylactic treatment.

[0052] In addition, the conditions, diseases, and maladies collectively referenced to as "Syndrome X" or Metabolic Syndrome as detailed in Johannsson *J. Clin. Endocrinol. Metab.*, 82, 727-34 (1997), may be treated employing the compounds of the invention.

25 B. Combinations

[0053] The present invention includes within its scope pharmaceutical compositions comprising, as an active ingredient, a therapeutically effective amount of at least one of the compounds of formula I, alone or in combination with a pharmaceutical carrier or diluent. Optionally, compounds of the present invention can be used alone, in combination with other compounds of the invention, or in combination with one or more other therapeutic agent(s), e.g., an antibiotic or other pharmaceutically active material.

[0054] The compounds of the present invention may be combined with growth promoting agents, such as, but not limited to, TRH, diethylstilbesterol, theophylline, enkephalins, E series prostaglandins, compounds disclosed in U.S. Patent No. 3,239,345, e.g., zeranol, and compounds disclosed in U.S. Patent No. 4,036,979, e.g., sulbenox or peptides disclosed in U.S. Patent No. 4,411,890.

[0055] The compounds of the invention may also be used in combination with growth hormone secretagogues such as GHRP-6, GHRP-1 (as described in U.S. Patent No. 4,411,890 and publications WO 89/07110 and WO 89/07111), GHRP-2 (as described in WO 93/04081), NN703 (Novo Nordisk), LY444711 (Lilly), MK-677 (Merck), CP424391 (Pfizer) and B-HT920, or with growth hormone releasing factor and its analogs or growth hormone and its analogs or somatomedins including IGF-1 and IGF-2, or with alpha-adrenergic agonists, such as clonidine or serotonin 5-HT_D agonists, such as sumatriptan, or agents which inhibit somatostatin or its release, such as physostigmine and pyridostigmine. A still further use of the disclosed compounds of the invention is in combination with parathyroid hormone, PTH(1-34) or bisphosphonates, such as MK-217 (alendronate).

[0056] A still further use of the compounds of the invention is in combination with estrogen, testosterone, a selective estrogen receptor modulator, such as tamoxifen or raloxifene, or other androgen receptor modulators, such as those disclosed in Edwards, J.P. *et. al.*, *Bio. Med. Chem. Let.*, 9, 1003-1008 (1999) and Hamann, L.G. *et. al.*, *J. Med. Chem.*, 42, 210-212 (1999).

[0057] A further use of the compounds of this invention is in combination with progesterone receptor agonists ("PRA"), such as levonorgestrel, medroxyprogesterone acetate (MPA).

[0058] The compounds of the present invention may be employed alone or in combination with each other and/or other modulators of nuclear hormone receptors or other suitable therapeutic agents useful in the treatment of the aforementioned disorders including: anti-diabetic agents; anti-osteoporosis agents; anti-obesity agents; anti-inflammatory agents; anti-anxiety agents; anti-depressants; anti-hypertensive agents; anti-platelet agents; anti-thrombotic and thrombolytic agents; cardiac glycosides; cholesterol/lipid lowering agents; mineralocorticoid receptor antagonists; phosphodiesterase inhibitors; protein tyrosine kinase inhibitors; thyroid

mimetics (including thyroid receptor agonists); anabolic agents; HIV or AIDS therapies; therapies useful in the treatment of Alzheimer's disease and other cognitive disorders; therapies useful in the treatment of sleeping disorders; anti-proliferative agents; and anti-tumor agents.

- 5 **[0059]** Examples of suitable anti-diabetic agents for use in combination with the compounds of the present invention include biguanides (e.g., metformin), glucosidase inhibitors (e.g., acarbose), insulins (including insulin secretagogues or insulin sensitizers), meglitinides (e.g., repaglinide), sulfonylureas (e.g., glimepiride, glyburide and glipizide), biguanide/glyburide combinations (e.g., Glucovance[®]),
- 10 thiazolidinediones (e.g., troglitazone, rosiglitazone and pioglitazone), PPAR-alpha agonists, PPAR-gamma agonists, PPAR alpha/gamma dual agonists, SGLT2 inhibitors, glycogen phosphorylase inhibitors, inhibitors of fatty acid binding protein (aP2) such as those disclosed in U.S. Serial No. 09/519,079 filed March 6, 2000, glucagon-like peptide-1 (GLP-1), and dipeptidyl peptidase IV (DPP4) inhibitors such
- 15 as those disclosed in WO 0168603.

- [0060]** Examples of suitable anti-osteoporosis agents for use in combination with the compounds of the present invention include alendronate, risedronate, PTH, PTH fragment, raloxifene, calcitonins, steroidal or non-steroidal progesterone receptor agonists, RANK ligand antagonists, calcium sensing receptor antagonists, TRAP
- 20 inhibitors, selective estrogen receptor modulators (SERM's), estrogen and AP-1 inhibitors.

- [0061]** Examples of suitable anti-obesity agents for use in combination with the compounds of the present invention include aP2 inhibitors, such as those disclosed in U.S. Serial No. 09/519,079 filed March 6, 2000, PPAR gamma antagonists, PPAR
- 25 delta agonists, beta 3 adrenergic agonists, such as AJ9677 (Takeda/Dainippon), L750355 (Merck), or CP331648 (Pfizer) or other known beta 3 agonists as disclosed in U.S. Patent Nos. 5,541,204, 5,770,615, 5,491,134, 5,776,983 and 5,488,064, a lipase inhibitor, such as orlistat or ATL-962 (Alizyme), a serotonin (and dopamine) reuptake inhibitor, such as sibutramine, topiramate (Johnson & Johnson) or axokine
- 30 (Regeneron), a thyroid receptor beta drug, such as a thyroid receptor ligand as disclosed in WO 97/21993 (U. Cal SF), WO 99/00353 (KaroBio) and GB98/284425

(KaroBio), and/or an anorectic agent, such as dexamphetamine, phentermine, phenylpropanolamine or mazindol.

[0062] Examples of suitable anti-inflammatory agents for use in combination with the compounds of the present invention include prednisone, dexamethasone, Enbrel[®],
5 cyclooxygenase inhibitors (i.e., COX-1 and/or COX-2 inhibitors such as NSAIDs, aspirin, indomethacin, ibuprofen, piroxicam, Naproxen[®], Celebrex[®], Vioxx[®]), CTLA4-Ig agonists/antagonists, CD40 ligand antagonists, IMPDH inhibitors, such as mycophenolate (CellCept[®]), integrin antagonists, alpha-4 beta-7 integrin antagonists, cell adhesion inhibitors, interferon gamma antagonists, ICAM-1, tumor necrosis
10 factor (TNF) antagonists (e.g., infliximab, OR1384), prostaglandin synthesis inhibitors, budesonide, clofazimine, CNI-1493, CD4 antagonists (e.g., priliximab), p38 mitogen-activated protein kinase inhibitors, protein tyrosine kinase (PTK) inhibitors, IKK inhibitors, and therapies for the treatment of irritable bowel syndrome (e.g., Zelmac[®] and Maxi-K[®] openers such as those disclosed in U.S. Patent No.
15 6,184,231 B1).

[0063] Examples of suitable anti-anxiety agents for use in combination with the compounds of the present invention include diazepam, lorazepam, buspirone, oxazepam, and hydroxyzine pamoate.

[0064] Examples of suitable anti-depressants for use in combination with the
20 compounds of the present invention include citalopram, fluoxetine, nefazodone, sertraline, and paroxetine.

[0065] Examples of suitable anti-hypertensive agents for use in combination with the compounds of the present invention include beta adrenergic blockers, calcium
25 channel blockers (L-type and T-type; e.g. diltiazem, verapamil, nifedipine, amlodipine and mybefradil), diuretics (e.g., chlorothiazide, hydrochlorothiazide, flumethiazide, hydroflumethiazide, bendroflumethiazide, methylchlorothiazide, trichloromethiazide, polythiazide, benzthiazide, ethacrynic acid, tricyclic, chlorthalidone, furosemide, musolimine, bumetanide, triamtrenene, amiloride, spironolactone), renin inhibitors, ACE inhibitors (e.g., captopril, zofenopril, fosinopril, enalapril, ceranopril, cilazopril,
30 delapril, pentopril, quinapril, ramipril, lisinopril), AT-1 receptor antagonists (e.g., losartan, irbesartan, valsartan), ET receptor antagonists (e.g., sitaxsentan, atrsentan

and compounds disclosed in U.S. Patent Nos. 5,612,359 and 6,043,265), Dual ET/AII antagonist (e.g., compounds disclosed in WO 00/01389), neutral endopeptidase (NEP) inhibitors, vasopepsidase inhibitors (dual NEP-ACE inhibitors) (e.g., omapatrilat and gemopatrilat), and nitrates.

- 5 [0066] Examples of suitable anti-platelet agents for use in combination with the compounds of the present invention include GPIIb/IIIa blockers (e.g., abciximab, eptifibatide, tirofiban), P2Y₁₂ antagonists (e.g., clopidogrel, ticlopidine, CS-747), thromboxane receptor antagonists (e.g., ifetroban), aspirin, and PDE-III inhibitors (e.g., dipyridamole) with or without aspirin.
- 10 [0067] Examples of suitable cardiac glycosides for use in combination with the compounds of the present invention include digitalis and ouabain.
- [0068] Examples of suitable cholesterol/lipid lowering agents for use in combination with the compounds of the present invention include HMG-CoA reductase inhibitors (e.g., pravastatin, lovastatin, atorvastatin, simvastatin, NK-104
- 15 (a.k.a. itavastatin, or nisvastatin or nisbastatin) and ZD-4522 (a.k.a. rosuvastatin, or atavastatin or visastatin)), squalene synthetase inhibitors, fibrates, bile acid sequestrants, ACAT inhibitors, MTP inhibitors, lipooxygenase inhibitors, cholesterol absorption inhibitors, and cholesterol ester transfer protein inhibitors (e.g., CP-529414).
- 20 [0069] Examples of suitable mineralocorticoid receptor antagonists for use in combination with the compounds of the present invention include spironolactone and eplerinone.
- [0070] Examples of suitable phosphodiesterase (PDE) inhibitors for use in combination with the compounds of the present invention include PDE-3 inhibitors
- 25 such as cilostazol, and phosphodiesterase-5 inhibitors (PDE-5 inhibitors) such as sildenafil.
- [0071] Examples of suitable thyroid mimetics for use in combination with the compounds of the present invention include thyrotropin, polythyroid, KB-130015, and dronedarone.
- 30 [0072] Examples of suitable anabolic agents for use in combination with the compounds of the present invention include testosterone, TRH diethylstilbesterol, estrogens, β -agonists, theophylline, anabolic steroids, dehydroepiandrosterone,

enkephalins, E-series prostagladins, retinoic acid and compounds as disclosed in U.S. Pat. No. 3,239,345, e.g., Zeranol[®]; U.S. Patent No. 4,036,979, e.g., Sulbenox[®] or peptides as disclosed in U.S. Pat. No. 4,411,890.

5 [0073] Examples of suitable HIV or AIDS therapies for use in combination with the compounds of the present invention include indinavir sulfate, saquinavir, saquinavir mesylate, ritonavir, lamivudine, zidovudine, lamivudine/zidovudine combinations, zalcitabine, didanosine, stavudine, and megestrol acetate.

[0074] Examples of suitable therapies for treatment of Alzheimer's disease and cognitive disorders for use in combination with the compounds of the present
10 invention include donepezil, tacrine, revastigmine, 5HT₆, gamma secretase inhibitors, beta secretase inhibitors, SK channel blockers, Maxi-K blockers, and KCNQs blockers.

[0075] Examples of suitable therapies for treatment of sleeping disorders for use in combination with the compounds of the present invention include melatonin
15 analogs, melatonin receptor antagonists, ML1B agonists, and GABA/NMDA receptor antagonists.

[0076] Examples of suitable anti-proliferative agents for use in combination with the compounds of the present invention include cyclosporin A, paclitaxel, FK-506, and adriamycin.

20 [0077] Examples of suitable anti-tumor agents for use in combination with the compounds of the present invention include paclitaxel, adriamycin, epothilones, cisplatin and carboplatin.

[0078] Compounds of the present invention may further be used in combination with nutritional supplements such as those described in U.S. 5,179,080, especially in
25 combination with whey protein or casein, amino acids (such as leucine, branched amino acids and hydroxymethylbutyrate), triglycerides, vitamins (e.g., A, B₆, B₁₂, folate, C, D and E), minerals (e.g., selenium, magnesium, zinc, chromium, calcium and potassium), carnitine, lipoic acid, creatinine, B-hydroxy-B-methylbutyrate (Juven) and coenzyme Q-10.

30 [0079] In addition, compounds of the present invention may be used in combination with therapeutic agents used in the treatment of sexual dysfunction, including but not limited to PDE-5 inhibitors, such as sildenafil or IC-351.

[0080] Compounds of the present invention may further be used in combination with antiresorptive agents, hormone replacement therapies, vitamin D analogues, elemental calcium and calcium supplements, cathepsin K inhibitors, MMP inhibitors, vitronectin receptor antagonists, Src SH₂ antagonists, vacuolar -H⁺- ATPase inhibitors, ipriflavone, fluoride, Tibolone, prostanoids, 17-beta hydroxysteroid dehydrogenase inhibitors and Src kinase inhibitors.

[0081] Compounds of the present invention may be used in combination with male contraceptives, such as nonoxynol 9 or therapeutic agents for the treatment of hair loss, such as minoxidil and finasteride or chemotherapeutic agents, such as with LHRH agonists.

[0082] Further, the compounds of the present invention may be used in combination with anti-cancer and cytotoxic agents, including but not limited to alkylating agents such as nitrogen mustards, alkyl sulfonates, nitrosoureas, ethylenimines, and triazenes; antimetabolites such as folate antagonists, purine analogues, and pyrimidine analogues; antibiotics such as anthracyclines, bleomycins, mitomycin, dactinomycin, and plicamycin; enzymes such as L-asparaginase; farnesyl-protein transferase inhibitors; 5 α -reductase inhibitors; inhibitors of 17 β -hydroxysteroid dehydrogenase type 3; hormonal agents such as glucocorticoids, estrogens/ antiestrogens, androgens/ antiandrogens, progestins, and luteinizing hormone-releasing hormone antagonists, octreotide acetate; microtubule-disruptor agents, such as ecteinascidins or their analogs and derivatives; microtubule-stabilizing agents such as taxanes, for example, paclitaxel (Taxol[®]), docetaxel (Taxotere[®]), and their analogs, and epothilones, such as epothilones A-F and their analogs; plant-derived products, such as vinca alkaloids, epipodophyllotoxins, taxanes; and topoisomerase inhibitors; prenyl-protein transferase inhibitors; and miscellaneous agents such as hydroxyurea, procarbazine, mitotane, hexamethylmelamine, platinum coordination complexes such as cisplatin and carboplatin; and other agents used as anti-cancer and cytotoxic agents such as biological response modifiers, growth factors; immune modulators and monoclonal antibodies. The compounds of the invention may also be used in conjunction with radiation therapy.

[0083] Representative examples of these classes of anti-cancer and cytotoxic agents include but are not limited to mechlorethamine hydrochloride,

cyclophosphamide, chlorambucil, melphalan, ifosfamide, busulfan, carmustin, lomustine, semustine, streptozocin, thiotepa, dacarbazine, methotrexate, thioguanine, mercaptopurine, fludarabine, pentastatin, cladribin, cytarabine, fluorouracil, doxorubicin hydrochloride, daunorubicin, idarubicin, bleomycin sulfate, mitomycin C, actinomycin D, safracins, saframycins, quinocarcins, discodermolides, vincristine, vinblastine, vinorelbine tartrate, etoposide, etoposide phosphate, teniposide, paclitaxel, tamoxifen, estramustine, estramustine phosphate sodium, flutamide, buserelin, leuprolide, pteridines, diynes, levamisole, aflacon, interferon, interleukins, aldesleukin, filgrastim, sargramostim, rituximab, BCG, tretinoin, irinotecan hydrochloride, betamethosone, gemcitabine hydrochloride, altretamine, and topotecan and any analogs or derivatives thereof.

[0084] Preferred member of these classes include, but are not limited to, paclitaxel, cisplatin, carboplatin, doxorubicin, carminomycin, daunorubicin, aminopterin, methotrexate, methopterin, mitomycin C, ecteinascidin 743, or porfiromycin, 5-fluorouracil, 6-mercaptopurine, gemcitabine, cytosine arabinoside, podophyllotoxin or podophyllotoxin derivatives such as etoposide, etoposide phosphate or teniposide, melphalan, vinblastine, vincristine, leurosine, vindesine and leurosine.

[0085] Examples of anticancer and other cytotoxic agents include the following: epothilone derivatives as found in German Patent No. 4138042.8; WO 97/19086, WO 98/22461, WO 98/25929, WO 98/38192, WO 99/01124, WO 99/02224, WO 99/02514, WO 99/03848, WO 99/07692, WO 99/27890, WO 99/28324, WO 99/43653, WO 99/54330, WO 99/54318, WO 99/54319, WO 99/65913, WO 99/67252, WO 99/67253 and WO 00/00485; cyclin dependent kinase inhibitors as found in WO 99/24416 (see also U.S. Patent No. 6,040,321); and prenyl-protein transferase inhibitors as found in WO 97/30992 and WO 98/54966; and agents such as those described generically and specifically in U.S. Patent No. 6,011,029 (the compounds of which U.S. Patent can be employed together with any NHR modulators (including, but not limited to, those of present invention) such as AR modulators, ER modulators, with LHRH modulators, or with surgical castration, especially in the treatment of cancer).

[0086] The above other therapeutic agents, when employed in combination with the compounds of the present invention, may be used, for example, in those amounts indicated in the Physicians' Desk Reference (PDR) or as otherwise determined by one of ordinary skill in the art.

5 [0087] The compounds of the formula I can be administered for any of the uses described herein by any suitable means, for example, orally, such as in the form of tablets, capsules, granules or powders; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intrasternal injection or infusion techniques (e.g., as sterile injectable aqueous or non-aqueous solutions or
10 suspensions); nasally, including administration to the nasal membranes, such as by inhalation spray; topically, such as in the form of a cream or ointment; or rectally such as in the form of suppositories; in dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents. The present compounds can, for example, be administered in a form suitable for immediate release or extended
15 release. Immediate release or extended release can be achieved by the use of suitable pharmaceutical compositions comprising the present compounds, or, particularly in the case of extended release, by the use of devices such as subcutaneous implants or osmotic pumps. The present compounds can also be administered liposomally.

[0088] Exemplary compositions for oral administration include suspensions
20 which can contain, for example, microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents such as those known in the art; and immediate release tablets which can contain, for example, microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and/or lactose and/or other
25 excipients, binders, extenders, disintegrants, diluents and lubricants such as those known in the art. The compounds of formula I can also be delivered through the oral cavity by sublingual and/or buccal administration. Molded tablets, compressed tablets or freeze-dried tablets are exemplary forms which may be used. Exemplary compositions include those formulating the present compound(s) with fast dissolving
30 diluents such as mannitol, lactose, sucrose and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (avicel) or polyethylene glycols (PEG). Such formulations can also include an excipient to aid

mucosal adhesion such as hydroxy propyl cellulose (HPC), hydroxy propyl methyl cellulose (HPMC), sodium carboxy methyl cellulose (SCMC), maleic anhydride copolymer (e.g., Gantrez), and agents to control release such as polyacrylic copolymer (e.g. Carbopol 934). Lubricants, glidants, flavors, coloring agents and stabilizers may also be added for ease of fabrication and use.

[0089] Exemplary compositions for nasal aerosol or inhalation administration include solutions in saline which can contain, for example, benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, and/or other solubilizing or dispersing agents such as those known in the art.

10 [0090] Exemplary compositions for parenteral administration include injectable solutions or suspensions which can contain, for example, suitable non-toxic, parenterally acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, or other suitable dispersing or wetting and suspending agents, including synthetic mono- or diglycerides, and fatty acids, including oleic acid, or Cremaphor.

15 [0091] Exemplary compositions for rectal administration include suppositories which can contain, for example, a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ordinary temperatures, but liquify and/or dissolve in the rectal cavity to release the drug.

20 [0092] Exemplary compositions for topical administration include a topical carrier such as Plastibase (mineral oil gelled with polyethylene).

[0093] The effective amount of a compound of the present invention can be determined by one of ordinary skill in the art, and includes exemplary dosage amounts for an adult human of from about 0.01 to 2000 mg of active compound per day, which can be administered in a single dose or in the form of individual divided doses, such as from 1 to 4 times per day. It will be understood that the specific dose level and frequency of dosage for any particular subject can be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the species, age, body weight, general health, sex and diet of the subject, the mode and time of administration, rate of excretion, drug combination, and severity of the particular condition. Preferred subjects for treatment include animals, most preferably

mammalian species such as humans, and domestic animals such as dogs, cats and the like, subject to NHR-associated conditions.

TRANSACTIVATION ASSAYS

5 AR Specific Assay

[0094] Compounds of the present invention were tested in transactivation assays of a transfected reporter construct and using the endogenous androgen receptor of the host cells. The transactivation assay provides a method for identifying functional agonists and partial agonists that mimic, or antagonists that inhibit, the effect of native
10 hormones, in this case, dihydrotestosterone (DHT). This assay can be used to predict *in vivo* activity as there is a good correlation in both series of data. See, e.g. T. Berger et al., *J. Steroid Biochem. Molec. Biol.* 773 (1992), the disclosure of which is herein incorporated by reference.

[0095] For the transactivation assay a reporter plasmid is introduced by
15 transfection (a procedure to induce cells to take foreign genes) into the respective cells. This reporter plasmid, comprising the cDNA for a reporter protein, such as secreted alkaline phosphatase (SEAP), controlled by prostate specific antigen (PSA) upstream sequences containing androgen response elements (AREs). This reporter plasmid functions as a reporter for the transcription-modulating activity of the AR.
20 Thus, the reporter acts as a surrogate for the products (mRNA then protein) normally expressed by a gene under control of the AR and its native hormone. In order to detect antagonists, the transactivation assay is carried out in the presence of constant concentration of the natural AR hormone (DHT) known to induce a defined reporter signal. Increasing concentrations of a suspected antagonist will decrease the reporter
25 signal (e.g., SEAP production). On the other hand, exposing the transfected cells to increasing concentrations of a suspected agonist will increase the production of the reporter signal.

[0096] For this assay, LNCaP and MDA 453 cells were obtained from the American Type Culture Collection (Rockville, MD), and maintained in RPMI 1640 or
30 DMEM medium supplemented with 10% fetal bovine serum (FBS; Gibco) respectively. The respective cells were transiently transfected by electroporation according to the optimized procedure described by Heiser, 130 Methods Mol. Biol.,

117 (2000), with the pSEAP2/PSA540/Enhancer reporter plasmid. The reporter plasmid, was constructed as follows: commercial human placental genomic DNA was used to generate by Polymerase Cycle Reaction (PCR) a fragment containing the BglII site (position 5284) and the Hind III site at position 5831 of the human prostate specific antigen promoter (Accession # U37672), Schuur, et al., *J. Biol. Chem.*, **271** (12): 7043-51 (1996). This fragment was subcloned into the pSEAP2/basic (Clontech) previously digested with BglII and HindIII to generate the pSEAP2/PSA540 construct. Then a fragment bearing the fragment of human PSA upstream sequence between positions -5322 and -3873 was amplified by PCR from human placental genomic DNA. A XhoI and a BglII sites were introduced with the primers. The resulting fragment was subcloned into pSEAP2/PSA540 digested with XhoI and BglII respectively, to generate the pSEAP2/PSA540/Enhancer construct. LNCaP and MDA MB-453 cells were collected in media containing 10% charcoal stripped FBS. Each cell suspension was distributed into two Gene Pulser Cuvetts (Bio-Rad) which then received 8 µg of the reporter construct, and electoporated using a Bio-Rad Gene Pulser at 210 volts and 960 µFaraday. Following the transfections the cells were washed and incubated with media containing charcoal stripped fetal bovine serum in the absence (blank) or presence (control) of 1 nM dihydrotestosterone (DHT; Sigma Chemical) and in the presence or absence of the standard anti-androgen bicalutamide or compounds of the present invention in concentrations ranging from 10^{-10} to 10^{-5} M (sample). Duplicates were used for each sample. The compound dilutions were performed on a Biomek 2000 laboratory workstation.

[0097] After 48 h, a fraction of the supernatant was assayed for SEAP activity using the Phospha-Light Chemiluminescent Reporter Gene Assay System (Tropix, Inc). Viability of the remaining cells was determined using the CellTiter 96 Aqueous Non-Radioactive Cell Proliferation Assay (MTS Assay, Promega). Briefly, a mix of a tetrazolium compound (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)- 2-(4-sulfophenyl)-2H-tetrazolium, inner salt; MTS) and an electron coupling reagent (phenazine methosulfate; PMS) are added to the cells. MTS (Owen's reagent) is bioreduced by cells into a formazan that is soluble in tissue culture medium, and therefore its absorbance at 490 nm can be measured directly from 96 well assay plates without additional processing. The quantity of formazan product as measured by the

amount of 490 nm absorbance is directly proportional to the number of living cells in culture. For each replicate the SEAP reading was normalized by the Abs490 value derived from the MTS assay. For the antagonist mode, the % Inhibition was calculated as:

5

$$\% \text{ Inhibition} = 100 \times (1 - [\text{average control} - \text{average blank} / \text{average sample} - \text{average blank}])$$

10 Data was plotted and the concentration of compound that inhibited 50% of the normalized SEAP was quantified (IC_{50}).

[0098] For the agonist mode % Control was referred as the effect of the tested compound compared to the maximal effect observed with the natural hormone, in this case DHT, and was calculated as:

15

$$\% \text{ Control} = 100 \times \text{average sample} - \text{average blank} / \text{average control} - \text{average blank}$$

Data was plotted and the concentration of compound that activates to levels 50% of the normalized SEAP for the control was quantified (EC_{50}).

20

GR Specificity Assay

[0099] The reporter plasmid utilized was comprised of the cDNA for the reporter SEAP protein, as described for the AR specific transactivation assay. Expression of the reporter SEAP protein was controlled by the mouse mammary tumor virus long terminal repeat (MMTV LTR) sequences that contains three hormone response elements (HREs) that can be regulated by both GR and PR see, e.g. G. Chalepakis et al., Cell, 53(3), 371 (1988). This plasmid was transfected into A549 cells, which expresses endogenous GR, to obtain a GR specific transactivation assay. A549 cells were obtained from the American Type Culture Collection (Rockville, MD), and maintained in RPMI 1640 supplemented with 10% fetal bovine serum (FBS; Gibco). Determination of the GR specific antagonist activity of the compounds of the present invention was identical to that described for the AR specific transactivation assay,

25
30

except that the DHT was replaced with 5 nM dexamethasone (Sigma Chemicals), a specific agonist for GR. Determination of the GR specific agonist activity of the compounds of the present invention was performed as described for the AR transactivation assay, wherein one measures the activation of the GR specific reporter system by the addition of a test compound, in the absence of a known GR specific agonists ligand.

PR Specific Assay

[00100] The reporter plasmid utilized was comprised of the cDNA for the reporter SEAP protein, as described for the AR specific transactivation assay. Expression of the reporter SEAP protein was controlled by the mouse mammary tumor virus long terminal repeat (MMTV LTR) sequences that contains three hormone response elements (HRE's) that can be regulated by both GR and PR. This plasmid was transfected into T47D, which expresses endogenous PR, to obtain a PR specific transactivation assay. T47D cells were obtained from the American Type Culture Collection (Rockville, MD), and maintained in DMEM medium supplemented with 10% fetal bovine serum (FBS; Gibco). Determination of the PR specific antagonist activity of the compounds of the present invention was identical to that described for the AR specific transactivation assay, except that the DHT was replaced with 1 nM Promegestone (NEN), a specific agonist for PR. Determination of the PR specific agonist activity of the compounds of the present invention was performed as described for the AR transactivation assay, wherein one measures the activation of the PR specific reporter system by the addition of a test compound, in the absence of a known PR specific agonists ligand.

AR Binding Assay

[00101] For the whole-cell binding assay, human LNCaP cells (T877A mutant AR) or MDA 453 (wild type AR) in 96-well microtiter plates containing RPMI 1640 or DMEM supplemented with 10% charcoal stripped CA-FBS (Cocaleco Biologicals) respectively, were incubated at 37 °C to remove any endogenous ligand that might be complexed with the receptor in the cells. After 48 h, either a saturation analysis to determine the K_d for tritiated dihydrotestosterone, [^3H]-DHT, or a competitive binding

assay to evaluate the ability of test compounds to compete with [³H]-DHT were performed. For the saturation analysis, media (RPMI 1640 or DMEM – 0.2% CA-FBS) containing [³H]-DHT (in concentrations ranging from 0.1 nM to 16 nM) in the absence (total binding) or presence (non-specific binding) of a 500-fold molar excess of unlabeled DHT were added to the cells. After 4 h at 37 °C, an aliquot of the total binding media at each concentration of [³H]-DHT was removed to estimate the amount of free [³H]-DHT. The remaining media was removed, cells were washed three times with PBS and harvested onto UniFilter GF/B plates (Packard), Microscint (Packard) was added and plates counted in a Top-Counter (Packard) to evaluate the amount of bound [³H]-DHT.

[00102] For the saturation analysis, the difference between the total binding and the non-specific binding, was defined as specific binding. The specific binding was evaluated by Scatchard analysis to determine the K_d for [³H]-DHT. See e.g. D. Rodbard, Mathematics and statistics of ligand assays: an illustrated guide: In: J. Langon and J. J. Clapp, eds., Ligand Assay, Masson Publishing U.S.A., Inc., New York, pp. 45-99, (1981), the disclosure of which is herein incorporated by reference.

[00103] For the competition studies, media containing 1 nM [³H]-DHT and compounds of the invention ("test compounds") in concentrations ranging from 10⁻¹⁰ to 10⁻⁵ M were added to the cells. Two replicates were used for each sample. After 4 h at 37 °C, cells were washed, harvested and counted as described above. The data was plotted as the amount of [³H]-DHT (% of control in the absence of test compound) remaining over the range of the dose response curve for a given compound. The concentration of test compound that inhibited 50% of the amount of [³H]-DHT bound in the absence of competing ligand was quantified (IC₅₀) after log-logit transformation. The K_i values were determined by application of the Cheng-Prusoff equation to the IC₅₀ values, where:

$$K_i = \frac{IC_{50}}{(1 + ([^3H\text{-DHT}] / K_d \text{ for } ^3H\text{-DHT}))}$$

30

After correcting for non-specific binding, IC₅₀ values were determined. The IC₅₀ is defined as the concentration of competing ligand needed to reduce specific binding by 50%. The K_ds for [³H]-DHT for MDA 453 and LNCaP were 0.7 and 0.2 nM respectively.

5

C2C12 Mouse Myoblast Transactivation Assay

[00104] Two functional transactivation assays were developed to assess the efficacy of androgen agonists in a muscle cell background using a luciferase reporter. The first assay (ARTA Stable 1) uses a cell line, Stable 1 (clone #72), which
10 expresses the full length rat androgen receptor but requires the transient transfection of an enhancer/reporter. This cell line was derived from C2C12 mouse myoblast cells. The second assay (ARTA Stable 2) uses a cell line, Stable 2 (clone #133), derived from Stable 1 which expresses both rAR and the enhancer/luciferase reporter.

[00105] The enhancer/reporter construct used in this system is pGL3/2XDR-
15 1/luciferase. 2XDR-1 was reported to be an AR specific response element in CV-1 cells, Brown et. al. The Journal of Biological Chemistry 272, 8227-8235, (1997). It was developed by random mutagenesis of an AR/GR consensus enhancer sequence.

ARTA Stable 1

- 20 1. Stable 1 cells are plated in 96 well format at 6,000 cells/well in high glucose DMEM without phenol red (Gibco BRL, Cat. No.: 21063-029) containing 10% charcoal and dextran treated FBS (HyClone Cat. No.: SH30068.02), 50 mM HEPES Buffer (Gibco BRL, Cat. No.: 15630-080), 1X MEM Na Pyruvate (Gibco BRL, Cat. No.: 11360-070), 0.5X Antibiotic-Antimycotic, and 800 µg/mL Geneticin
25 (Gibco BRL, Cat. No.: 10131-035).
2. 48 h later, cells are transfected with pGL3/2XDR-1/luciferase using LipofectAMINE PlusTM Reagent (Gibco BRL, Cat. No.: 10964-013). Specifically, 5 ng/well pGL3/2XDR-1/luciferase DNA and 50 ng/well Salmon Sperm DNA (as
30 070). To this, 0.5 µl/well Plus reagent is added. This mixture is incubated for 15 min at rt. In a separate vessel, 0.385 µl/well LipofectAMINE reagent is diluted with 5

μl/well Opti-MEM. The DNA mixture is then combined with the LipofectAMINE mixture and incubated for an additional 15 min at rt. During this time, the media from the cells is removed and replaced with 60 μl/well of Opti-MEM. To this is added 10 μl/well of the DNA/LipofectAMINE transfection mixture. The cells are incubated for 4 h.

3. The transfection mixture is removed from the cells and replaced with 90 μl of media as in #1 above.

4. 10 μl/well of appropriate drug dilution is placed in each well.

5. 24 h later, the Steady-GloTM Luciferase Assay System is used to detect activity according to the manufacturer's instructions (Promega, Cat. No.: E2520).

ARTA Stable 2

1. Stable 2 cells are plated in 96 well format at 6,000 cells/well in high glucose DMEM without phenol red (Gibco BRL, Cat. No.: 21063-029) containing 10% charcoal and dextran treated FBS (HyClone Cat. No.: SH30068.02), 50 mM HEPES Buffer (Gibco BRL, Cat. No.: 15630-080), 1X MEM Na Pyruvate (Gibco BRL, Cat. No.: 11360-070), 0.5X Antibiotic-Antimycotic, 800 μg/mL Geneticin (Gibco BRL, Cat. No.: 10131-035) and 800 μg/mL Hygromycin β (Gibco BRL, Cat. No.: 10687-010).

2. 48 h later, the media on the cells is removed and replaced with 90 μl fresh. 10 μl/well of appropriate drug dilution is placed in each well.

3. 24 h later, the Steady-GloTM Luciferase Assay System is used to detect activity according to the manufacturer's instructions (Promega, Cat. No. E2520).

PROLIFERATION ASSAYS

Human Prostate Cell Proliferation Assay

[00106] Compounds of the present invention were tested ("test compounds") on the proliferation of human prostate cancer cell lines. For that, MDA PCa2b cells, a cell line derived from the metastasis of a patient that failed castration, Navone et al., Clin. Cancer Res., 3, 2493-500 (1997), were incubated with or without the test compounds

for 72 h and the amount of [³H]-thymidine incorporated into DNA was quantified as a way to assess number of cells and therefore proliferation. The MDA PCa2b cell line was maintained in BRFF-HPC1 media (Biological Research Faculty & Facility Inc., MD) supplemented with 10% FBS. For the assay, cells were plated in Biocoated 96-well microplates and incubated at 37 °C in 10% FBS (charcoal-stripped)/BRFF-BMZERO (without androgens). After 24 h, the cells were treated in the absence (blank) or presence of 1 nM DHT (control) or with test compounds (sample) of the present invention in concentrations ranging from 10⁻¹⁰ to 10⁻⁵ M. Duplicates were used for each sample. The compound dilutions were performed on a Biomek 2000 laboratory work station. Seventy-two h later 0.44 uCi. of [³H]-Thymidine (Amersham) was added per well and incubated for another 24 h followed by trypsinization, harvesting of the cells onto GF/B filters. Micro-scint PS were added to the filters before counting them on a Beckman TopCount.

[00107] The % Inhibition was calculated as:

$$\% \text{ Inhibition} = 100 \times (1 - [\text{average}_{\text{control}} - \text{average}_{\text{blank}} / \text{average}_{\text{sample}} - \text{average}_{\text{blank}}])$$

Data was plotted and the concentration of compound that inhibited 50% of the [³H]-Thymidine incorporation was quantified (IC₅₀).

Murine Breast Cell Proliferation Assay

[00108] The ability of compounds of the present invention ("test compounds") to modulate the function of the AR was determined by testing said compounds in a proliferation assay using the androgen responsive murine breast cell line derived from the Shionogi tumor, Hiraoka *et al.*, *Cancer Res.*, **47**, 6560-6564 (1987). Stable AR dependent clones of the parental Shionogi line were established by passing tumor fragments under the general procedures originally described in Tetuo, *et. al.*, *Cancer Research* **25**, 1168-1175 (1965). From the above procedure, one stable line, SC114, was isolated, characterized and utilized for the testing of example compounds. SC114 cells were incubated with or without the test compounds for 72 h and the amount of [³H]-thymidine incorporated into DNA was quantified as a surrogate endpoint to

assess the number of cells and therefore the proliferation rate as described in Suzuki *et. al.*, *J. Steroid Biochem. Mol. Biol.* 37, 559-567 (1990). The SC114 cell line was maintained in MEM containing 10^{-8} M testosterone and 2% DCC-treated FCS. For the assay, cells were plated in 96-well microplates in the maintenance media and
5 incubated at 37 °C. On the following day, the medium was changed to serum free medium [Ham's F-12:MEM (1;1, v/v) containing 0.1% BSA] with (antagonist mode) or without (agonist mode) 10^{-8} M testosterone and the test compounds of the present invention in concentrations ranging from 10^{-10} to 10^{-5} M. Duplicates were used for each sample. The compound dilutions were performed on a Biomek 2000 laboratory
10 work station. Seventy two h later 0.44uCi of [3H]-Thymidine (Amersham) was added per well and incubated for another 2 h followed by trypsinization, and harvesting of the cells onto GF/B filters. Micro-scint PS were added to the filters before counting them on a Beckman TopCount.

15 [00109] For the antagonist mode, the % Inhibition was calculated as:

$$\% \text{ Inhibition} = 100 \times (1 - [\text{average}_{\text{sample}} - \text{average}_{\text{blank}} / \text{average}_{\text{control}} - \text{average}_{\text{blank}}])$$

Data was plotted and the concentration of compound that inhibited 50% of the [3 H]-
20 Thymidine incorporation was quantified (IC_{50}).

[00110] For the agonist mode % Control was referred as the effect of the tested compound compared to the maximal effect observed with the natural hormone, in this case DHT, and was calculated as:

25

$$\% \text{ Control} = 100 \times (\text{average}_{\text{sample}} - \text{average}_{\text{blank}}) / (\text{average}_{\text{control}} - \text{average}_{\text{blank}})$$

Data was plotted and the concentration of compound that inhibited 50% of the [3 H]-
Thymidine incorporation was quantified (EC_{50}).

30

***In Vitro* Assay to Measure GR-Induced AP-1 Transrepression**

[00111] The AP-1 assay is a cell-based luciferase reporter assay. A549 cells, which contain endogenous glucocorticoid receptor, were stably transfected with an AP-1 DNA binding site attached to the luciferase gene. Cells are then grown in RPMI + 10% fetal calf serum (charcoal-treated) + Penicillin/Streptomycin with 0.5mg/mL geneticin. Cells are plated the day before the assay at approximately 40000 cells/well. On assay day, the media is removed by aspiration and 20 μ L assay buffer (RPMI without phenol red + 10% FCS (charcoal-treated) + Pen/Strep) is added to each well. At this point either 20 μ L assay buffer (control experiments), the compounds of the present invention ("test compounds") (dissolved in DMSO and added at varying concentrations) or dexamethasone (100 nM in DMSO, positive control) are added to each well. The plates are then pre-incubated for 15 min at 37 °C, followed by stimulation of the cells with 10 ng/mL PMA. The plates are then incubated for 7 h at 37 °C after which 40 μ L luciferase substrate reagent is added to each well. Activity is measured by analysis in a luminometer as compared to control experiments treated with buffer or dexamethasone. Activity is designated as % inhibition of the reporter system as compared to the buffer control with 10 ng/mL PMA alone. The control, dexamethasone, at a concentration of ≤ 10 μ M typically suppresses activity by 65%. Test compounds which demonstrate an inhibition of PMA induction of 50% or greater at a concentration of test compound of ≤ 10 μ M are deemed active.

IN VIVO ASSAYS**Levator Ani & Wet Prostate Weight Assay AR Agonist Assay**

[00112] The activity of compounds of the present invention as AR agonists was investigated in an immature male rat model, a recognized test of anabolic effects in muscle and sustaining effects in sex organs for a given compound, as described in L. G. Hershberger et al., *Proc. Soc. Expt. Biol. Med.*, **83**, 175 (1953); B. L. Beyler et al, "Methods for evaluating anabolic and catabolic agents in laboratory animals", *J. Amer. Med. Women's Ass.*, **23**, 708 (1968); H. Fukuda et al., "Investigations of the levator ani muscle as an anabolic steroid assay", *Nago Dai. Yak. Ken. Nem.* **14**, 84 (1966) the disclosures of which are herein incorporated by reference.

[00113] The basis of this assay lies in the well-defined action of androgenic agents on the maintenance and growth of muscle tissues and sexual accessory organs in animals and man. Androgenic steroids, such as testosterone (T), have been well characterized for their ability to maintain muscle mass. Treatment of animals or humans after castrations with an exogenous source of T results in a reversal of muscular atrophy. The effects of T on muscular atrophy in the rat levator ani muscle have been well characterized. M. Masuoka et al., "Constant cell population in normal, testosterone deprived and testosterone stimulated levator ani muscles" *Am. J. Anat.* **119**, 263 (1966); Z. Gori et al., "Testosterone hypertrophy of levator ani muscle of castrated rats. I. Quantitative data" *Boll. -Soc. Ital. Biol. Sper.* **42**, 1596 (1966); Z. Gori et al., "Testosterone hypertrophy of levator ani muscle of castrated rats. II. Electron-microscopic observations" *Boll. -Soc. Ital. Biol. Sper.* **42**, 1600 (1966); A. Boris et al., *Steroids* **15**, 61 (1970). As described above, the effects of androgens on maintenance of male sexual accessory organs, such as the prostate and seminal vesicles, is well described. Castration results in rapid involution and atrophy of the prostate and seminal vesicles. This effect can be reversed by exogenous addition of androgens. Since both the levator ani muscle and the male sex organs are the tissues most responsive to the effects of androgenic agents, this model is used to determine the androgen dependent reversal of atrophy in the levator ani muscle and the sex accessory organs in immature castrated rats. Sexually mature rats (200-250 g, 6-8 weeks-old, Sprague-Dawley, Harlan) were acquired castrated from the vendor (Taconic). The rats were divided into groups and treated daily for 7 to 14 days with one of the following:

1. Control vehicle
2. Testosterone Propionate (TP) (3 mg/rat/day, subcutaneous)
3. TP plus Bicalutamide (administered p.o. in PEGTW, QD), a recognized antiandrogen, as a reference compound.
4. To demonstrate antagonist activity, a compound of the present invention ("test compound") was administered (p.o. in PEGTW, QD) with TP (s.c. as administered in group 2) in a range of doses.
5. To demonstrate agonist activity a compound of the present invention ("test compound") was administered alone (p.o. in PEGTW, QD) in a range of doses.

[00114] At the end of the 7-14-day treatment, the animals were sacrificed by carbon dioxide, and the levator ani, seminal vesicle and ventral prostate weighed. To compare data from different experiments, the levator ani muscle and sexual organ weights were first standardized as mg per 100 g of body weight, and the increase in organ weight induced by TP was considered as the maximum increase (100%).
5 Super-anova (one factor) was used for statistical analysis.

[00115] The gain and loss of sexual organ weight reflect the changes of the cell number (DNA content) and cell mass (protein content), depending upon the serum androgen concentration. See Y. Okuda et al., *J. Urol.*, **145**, 188-191 (1991), the
10 disclosure of which is herein incorporated by reference. Therefore, measurement of organ wet weight is sufficient to indicate the bioactivity of androgens and androgen antagonist. In immature castrated rats, replacement of exogenous androgens increases levator ani, seminal vesicles (SV) and prostate in a dose dependent manner.

[00116] The maximum increase in organ weight was 4 to 5-fold when dosing 3
15 mg/rat/day of testosterone (T) or 1 mg/rat/day of testosterone propionate (TP) for 3 days. The EC₅₀ of T and TP were about 1 mg and 0.03 mg, respectively. The increase in the weight of the VP and SV also correlated with the increase in the serum T and DHT concentration. Although administration of T showed 5-times higher serum concentrations of T and DHT at 2 h after subcutaneous injection than that of
20 TP, thereafter, these high levels declined very rapidly. In contrast, the serum concentrations of T and DHT in TP-treated animals were fairly consistent during the 24 h, and therefore, TP showed about 10-30-fold higher potency than free T.


EXAMPLES

[00117] The following examples serve to better illustrate, but not limit, some of the preferred embodiments of the invention.


5

EXAMPLE 1


2-Chloro-4-((7*S*,7*aR*)-7-hydroxy-1-methyl-3-oxo-5,6,7,7*a*-tetrahydro-3*H*-pyrrolizin-2-yl)-3-methylbenzonitrile



10

1A. 3-Chloro-2-methylphenylacetamide


- 15 [00118] To a solution of 3-chloro-2-methylaniline (3.00 g, 21.2 mmol) in 25 mL of EtOH at rt was added acetic anhydride (2.40 mL, 25.4 mmol), and the solution was stirred at rt for 2 h. The mixture was concentrated under reduced pressure to give 3.89 g (100%) of the desired acetamide. ¹H NMR (DMSO-*d*₆) δ 2.05 (s, 3H), 2.20 (s, 3H), 7.16 (t, *J* = 7.7, 8.3, 1H), 7.25 (d, *J* = 8.3, 1H), 7.31 (d, *J* = 8.3, 1H), 9.55 (s, 1H);
- 20 ¹³C NMR (DMSO-*d*₆) δ 15.1, 23.1, 124.4, 125.8, 126.7, 130.3, 133.7, 138.0, 168.3; HPLC a) column: Phenominex ODS C18 4.6 x 50 mm, 4 min gradient, 10% MeOH/90% H₂O/0.1% TFA to 90% MeOH/10% H₂O/0.1% TFA; 1 min hold, 4 mL/min UV detection at 220 nm, 2.32 min retention time; HPLC b) column: Shimadzu Shim-Pack VP-ODS C18 4.6 x 50 mm, 4 min gradient, 10% MeOH/90% H₂O/0.1% TFA to 90% MeOH/10% H₂O/0.1% TFA, 1 min hold; 4 mL/min, UV
- 25 detection at 220 nm, 2.20 min retention time (99%); MS (ES) *m/z* 184 [M+H]⁺.

1B. 4-Bromo-3-chloro-2-methylphenylacetamide

5 [00119] To a suspension of acetamide **1A** (2.00 g, 10.9 mmol) in 15 mL of glacial AcOH cooled to approximately 15 °C was added bromine (1.67 mL, 32.7 mmol) over 20 min. The ice bath was removed and the solution was stirred for 2 h, poured into ice water with stirring, and the solid was then filtered and dried to give 2.75 g (96%) of the desired bromide. ¹H NMR (DMSO-*d*₆) δ 2.05 (s, 3H), 2.28 (s, 3H), 7.29 (d, *J* = 8.3, 1H), 7.56 (d, *J* = 8.8, 1H), 9.60 (s, 1H); ¹³C NMR (DMSO-*d*₆) δ 16.7, 23.1, 118.1, 125.5, 130.4, 132.7, 133.4, 137.1, 168.4; HPLC a) column: Phenominex ODS C18 4.6 x 50 mm, 4 min gradient, 10% MeOH/90% H₂O/0.1% TFA to 90% MeOH/10% H₂O/0.1% TFA, 1 min hold, 4 mL/min, UV detection at 220 nm, 2.95 min retention time; HPLC b) column: Shimadzu Shim-Pack VP-ODS C18 4.6 x 50 mm, 4 min gradient, 10% MeOH/90% H₂O/0.1% TFA to 90% MeOH/10% H₂O/0.1% TFA, 1 min hold, 4 mL/min, UV detection at 220 nm, 2.87 min retention time (98%); MS (ES) *m/z* 263 [M+H]⁺.

1C. 3-Chloro-4-cyano-2-methylphenylacetamide


20




[00120] A suspension of bromide **1B** (2.70 g, 10.3 mmol) and copper cyanide (0.92 g, 10.3 mmol) in DMF (30 mL) was heated to 150 °C for 4 h. The suspension was cooled, poured into water with stirring, and the solid was filtered and dried to give 1.44 g (67%) of the desired nitrile. ¹H NMR (DMSO-*d*₆) δ 2.12 (s, 3H), 2.29 (s, 3H), 7.72 (d, *J* = 8.8, 1H), 7.75 (d, *J* = 8.2, 1H), 9.73 (s, 1H); ¹³C NMR (DMSO-*d*₆) δ 15.3,

23.5, 107.7, 116.5, 123.0, 130.1, 131.5, 135.7, 142.3, 168.8; HPLC a) column: Phenominex ODS C18 4.6 x 50 mm, 4 min gradient, 10% MeOH/90% H₂O/0.1% TFA to 90% MeOH/10% H₂O/0.1% TFA, 1 min hold, 4 mL/min, UV detection at 220 nm, 2.23 min retention time; HPLC b) column: Shimadzu Shim-Pack VP-ODS C18 4.6 x 50 mm, 4 min gradient, 10% MeOH/90% H₂O/0.1% TFA to 90% MeOH/10% H₂O/0.1% TFA, 1 min hold, 4 mL/min, UV detection at 220 nm, 2.13 min retention time (95%); MS (ES) *m/z* 209 [M+H]⁺.

1D. 3-Chloro-4-cyano-2-methylphenylaniline




[00121] A solution of cyanoacetamide **1C** (9.90 g, 47.4 mmol) in 100 mL of concentrated HCl/EtOH (1:1) was refluxed 30 min. The solution was then concentrated and dried under reduced pressure to give 9.41 g (98%) of the desired aniline as the hydrochloride salt. The free base of the aniline was obtained by suspending the salt in EtOAc and washing with saturated aqueous NaHCO₃ solution. The organic layer was then dried (MgSO₄), filtered and concentrated under reduced pressure. ¹H NMR (DMSO-*d*₆) δ 2.12 (s, 3H), 6.30 (s, 2H), 6.61 (d, *J* = 8.23, 1H), 7.36 (d, *J* = 8.23, 1H); ¹³C NMR (DMSO-*d*₆) δ 13.8, 96.9, 112.1, 118.3, 118.85, 132.2, 135.6, 152.5; HPLC a) column: Phenominex ODS C18 4.6 x 50 mm, 4 min gradient, 10% MeOH/90% H₂O/0.1% TFA to 90% MeOH/10% H₂O/0.1% TFA, 1 min hold, 4 mL/min, UV detection at 220 nm, 2.43 min retention time; HPLC b); column: Shimadzu Shim-Pack VP-ODS C18 4.6 x 50 mm, 4 min gradient, 10% MeOH/90% H₂O/0.1% TFA to 90% MeOH/10% H₂O/0.1% TFA, 1 min hold, 4 mL/min, UV detection at 220 nm, 2.31 min retention time (99%); MS (ES) *m/z* 167 [M+H]⁺.

1E. 2-Chloro-4-Iodo-3-methylbenzonitrile

5 [00122] To a suspension of CuI (7.5 g, 39.3 mmol) in acetonitrile (150 mL) under N₂ at rt was added *tert*-butylnitrite (5.7 mL, 47.9 mmol). The reaction mixture was heated to 65 °C for 1 h and then compound 1D (6.0 g, 36.0 mmol) was added and the reaction was heated at 65 °C for 3 h. The reaction was cooled to rt and filtered through a pad of celite. The celite pad was washed with EtOAc. The organics were
10 washed twice with water, dried over MgSO₄, filtered and concentrated. Purification by flash chromatography (silica gel, 110 g ISCO, 0-5% EtOAc in hexane, step gradient) gave the title compound (4.3 g): MS (ES) *m/z* 278 [M+H]⁺.


1F. Diethyl 2-(3-chloro-4-cyano-2-methylphenyl)malonate

15




[00123] To a pressure reaction flask containing diethyl malonate (4.1 mL, 27 mmol) in 50 mL of anhydrous dioxane was added NaH (1.1 g, 27.5 mmol, ~60% in mineral oil) in several portions at RT, and stirred for 30 min. Compound 1E (2.5 g, 9
20 mmol) and CuI (5.1 g, 26.8 mmol) were added, sealed and heated at 100 °C for 3 days. The reaction was cooled to RT and filtered through a pad of celite. The celite pad was washed with EtOAc. The organics were washed twice with water, dried over Na₂SO₄, filtered and concentrated. Purification by flash chromatography (silica gel,
25 110 g ISCO, 10% EtOAc in hexane) gave the title compound (2.3 g).

1G. 2-(3-Chloro-4-cyano-2-methylphenyl)acetic acid



5 **[00124]** To compound 1F (2.3 g, 7.4 mmol) in 20 mL of MeOH was added 1N
NaOH solution (30 mL, 30 mmol), and stirred at RT for 5 hr. It was concentrated,
EtOAc was added, then acidified with conc. HCl. The biphasic solution was heated at
60 °C for 2 hr, and cooled to RT. The organic layer was separated, dried over
Na₂SO₄, filtered and concentrated to give the title compound (1.44 g). MS (ES) *m/z*
10 164 [M-CO₂H]⁻.


1H. (2S,3S)- N-tert-butyloxycarbonyl-3-(tert-butyldimethylsilyloxy)-proline methyl ester



15


[00125] Following the procedures described in WO2003096980, (2*S*,3*S*)-3-hydroxyproline was converted to the title compound.

20 **1I. (2*R*,3*S*)-*N*-*tert*-Butyloxycarbonyl-3-(*tert*-butyldimethylsilanyloxy)-2-hydroxymethylpyrrolidine**




[00126] To a solution of compound **1H** (4.3 g, 12 mmol) in THF (75 mL) at -78°C under nitrogen was added dropwise 1.0 M lithium triethylborohydride in THF (60 mL, 60 mmol). The reaction was allowed to warm to rt and stirred for 4 h. The reaction was quenched by pouring over ice (150 g) and stirring for 30 min. The product was
5 extracted into EtOAc and washed with saturated aqueous NaHCO_3 . The organics were dried over MgSO_4 , filtered and concentrated. Purification by flash chromatography (silica gel, 0-15% EtOAc in hexanes) provided the title compound (3.3 g).

10 **1J. (2S,3S)-N-tert-Butyloxycarbonyl-3-(tert-butyldimethylsilyloxy)-2-formyl-pyrrolidine**



15 [00127] To a solution of compound **1I** (3.3 g, 9.97 mmol) in 10% CH_3CN in CH_2Cl_2 (50 mL) at rt was added NMO (3.0g, 25.6 mmol) followed by tetrapropyl ammonium perruthenate (300 mg, 0.85 mmol). After a mild exotherm, the reaction was stirred at rt for 4 h. The reaction was diluted with hexane (50 mL), mixed well
20 and filtered through a plug of silica gel (~50 g). The silica plug was washed liberally with 30% EtOAc in hexane (500 mL). Solvents were removed under vacuum. Purification by flash chromatography (silica gel, 0-15% EtOAc in hexanes) provided the title compound (2.94 g).

1K. (2R,3S)-N-tert-Butyloxycarbonyl-3-(tert-butyldimethylsilyloxy)-2-(1-hydroxyethyl)pyrrolidine




5

[00128] To a solution of **1J** (~0.05 M in THF) at -78 °C was added dropwise MeMgBr (3.0M in Et₂O, 3 equiv). The reaction was stirred at -78 °C for 1.5-2 h (quench small aliquot in 10% MeOH/DCM, spot TLC -silica gel, 30% EtOAc/Hex – stain with 5% HCl/MeOH, heat and then ninhydrin, heat). Quench reaction by addition of HOAc (3 equiv) at -78 °C. Warm to rt, dilute with EtOAc and wash with a 1:1 mixture of saturated aq. NaHCO₃ and brine. Back extract aqueous layer. Combine organic extracts, dry over MgSO₄, filter and concentrate. Purification by flash chromatography (silica gel, step gradient 0-10-20% EtOAc in hexane) provided the title compound.

15

1L. (2S,3S)-N-tert-Butyloxycarbonyl-3-(tert-butyldimethylsilyloxy)-2-(1-hydroxyethyl) pyrrolidine




20

[00129] To compound **1K** (0.1M solution in 10% CH₃CN in DCM) add NMO (2 equiv) followed by TPAP (0.05 equiv). Provide sufficient ventilation and take care to watch for exotherm. Stir reaction at rt for 3-5 h. Dilute by half with hexane and allow TPAP to precipitate. Flash filter through silica gel plug (20 fold mass of


starting alcohol) and elute with 10% EtOAc/Hex until product is not observed in eluent. Removal of solvents gave the title compound. MS (ES) m/z 344.31 $[M+H]^+$.

5 **1M. 4-(2-((2*S*,3*S*)-2-Acetyl-3-(tert-butyldimethylsilyloxy)pyrrolidin-1-yl)-2-oxoethyl)-2-chloro-3-methylbenzonitrile**



[00130] To compound 1L (480 mg, 1.4 mmol) was added 7 mL of 10%
10 TFA/CH₂Cl₂ solution at RT, and stirred at 4 hr. A 7 mL toluene was added to the reaction mixture, and concentrated. It was dissolved in 5 mL of CH₂Cl₂, a solution of PyBrop (1.2 g, 2.5 mmol) and compound 1G (210 mg, 1 mmol) in 3 mL of THF were added followed by iPr₂NET (520 μ L, 3 mmol), and the stirring was continued for 3 hr at RT. It was concentrated, dissolved in EtOAc, washed with water, dried
15 over Na₂SO₄, and concentrated. Purification by flash chromatography (silica gel, 30% EtOAc in hexanes) provided the title compound (220 mg). MS (ES) m/z 435 $[M+H]^+$.


20 **1N. 4-((7*S*)-7-(tert-butyldimethylsilyloxy)-1-hydroxy-1-methyl-3-oxo-hexahydro-1H-pyrrolizin-2-yl)-2-chloro-3-methylbenzonitrile**



[00131] To a pressure reaction vessel containing compound 1M (210 mg, 0.48 mmol) in 8 mL of anhydrous THF was added piperidine (240 μ L, 2.4 mmol). The


vessel was sealed, heated at 100 °C for 15 hr, and cooled to RT. Purification by flash chromatography (silica gel, 10-30% EtOAc in hexanes) provided the title compound (87 mg). MS (ES) m/z 435 $[M+H]^+$.

5 **10. 4-((7*S*)-7-(tert-butyldimethylsilyloxy)-1-methyl-3-oxo-5,6,7,7a-tetrahydro-3H-pyrrolizin-2-yl)-2-chloro-3-methylbenzonitrile**



10 **[00132]** To a solution of compound 1N (97 mg, 0.22 mmol) in 1.5 mL of anhydrous THF were added pyridine (180 μ L, 2.2 mmol) and TFAA (157 μ L, 1.1 mmol) at 0 °C. Ice-water bath was removed, the stirring was continued for 5 hr at RT, then quenched with water. It was concentrated, redissolved in EtOAc, washed with water, dried over Na_2SO_4 , and concentrated. The residue was dissolved in 1 mL of
15 anhydrous THF, $i\text{Pr}_2\text{NEt}$ (115 μ L, 0.66 mmol) was added. The reaction mixture was heated at 50 °C for 3 hr, cooled to RT, and concentrated. Purification by flash chromatography (silica gel, 10-30% EtOAc in hexanes) provided the title compound (42 mg). MS (ES) m/z 417 $[M+H]^+$.

20 **2-Chloro-4-((7*S*,7*aR*)-7-hydroxy-1-methyl-3-oxo-5,6,7,7a-tetrahydro-3H-pyrrolizin-2-yl)-3-methylbenzonitrile**




25 **[00133]** To a plastic vial containing compound 1O (32 mg, 0.076 mmol) in 1 mL of THF was added 100 μ L of HF pyridine solution (Aldrich), and stirred at RT


overnight. It was concentrated and purified on prepHPLC to give the title compound (17.5 mg). MS (ES) m/z 305 $[M+H]^+$.

EXAMPLE 2


5 4-((7*S*,7*aR*)-7-Hydroxy-1-methyl-3-oxo-5,6,7,7*a*-tetrahydro-3*H*-pyrrolizin-2-yl)-1-naphthonitrile



10 2A. (4-Cyanonaphthalen-1-yl)acetic acid methyl ester




[00134] To a solution of 4-methyl-1-cyanonaphthalene (4.35 g, 25.9 mmol) in THF
15 (100 mL) was cooled to -78°C and treated with commercially available LDA (2 M, 60 mL). The reaction mixture was then warmed to 0°C and maintained at 0°C for 30 min. The resulting red-colored solution was cooled to -78°C and methyl chloroformate (10 mL, 130 mmol) was added. The reaction mixture was stirred while warming to rt until the displacement was complete (4 h) and then quenched with sat'd
20 NH_4Cl . The THF layer was evaporated and extracted with EtOAc (3x20 mL). The EtOAc was dried over Na_2SO_4 and concentrated *in vacuo* to afford an oil.

2B. (4-Cyanonaphthalen-1-yl)acetic acid

5 [00135] To a solution of compound 2A (6.20 g, 26.7 mmol) in MeOH (100 mL) was treated with saturated aqueous LiOH (15 mL), and the reaction mixture was stirred at rt overnight (~16 h). The resulting mixture was concentrated *in vacuo* to dryness and partitioned between a mixed solvent (50% Et₂O:50% EtOAc) and H₂O. The H₂O layer was acidified to pH 2-3 and extracted with EtOAc (3x20 mL). The
10 EtOAc was dried over Na₂SO₄ and concentrated *in vacuo* to give a yellow solid (2.1 g) in 38% yield.

4-((7*S*,7*aR*)-7-Hydroxy-1-methyl-3-oxo-5,6,7,7*a*-tetrahydro-3*H*-pyrrolizin-2-yl)-1-naphthonitrile


15




[00136] The title compound was prepared from compound 1L and compound 2B in a similar manner as described in Example 1.

20

EXAMPLE 3

2-Chloro-4-((1*R*,7*S*,7*aS*)-1,7-dihydroxy-3-oxo-1-(trifluoromethyl)-hexahydro-1*H*-pyrrolizin-2-yl)-3-methylbenzonitrile


5

3A. (2*S*,3*S*)-Methyl 3-(tert-butyldimethylsilyloxy)-1-(2-(3-chloro-4-cyano-2-methylphenyl)acetyl)pyrrolidine-2-carboxylate

10

[00137] The title compound 3A was prepared from compound 1G and 1H in a similar manner as for compound 1M in Example 1. MS (ES) m/z 337 $[M+H]^+$ and 359 $[M+Na]^+$.


15

3B. 2-Chloro 4-((1*R*,7*S*,7*aS*)-7-(tert-butyldimethylsilyloxy)-1-hydroxy-3-oxo-1-(trifluoromethyl)-hexahydro-1*H*-pyrrolizin-2-yl)-3-methyl-benzonitrile

20

[00138] To compound 3A (970 mg, 2.2 mmol) in THF (1 mL) was added TMSiCF₃ (0.95 mL, 6.4 mmol) at RT followed by dry CsF (60 mg, 0.39 mmol; dried at 140 °C under vacuum). The reaction mixture was stirred at RT overnight, then diluted with 4N HCl/dioxane (2 mL) and MeOH (2 mL), and the stirring was continued for 2 hr at RT. It was concentrated, dissolved in EtOAc, washed with water, dried over Na₂SO₄, and concentrated to give a yellow residue. The residue was dissolved in 5 mL of CH₂Cl₂, imidazole (730 mg, 10.7 mmol) and TBSiCl (960 mg, 6.4 mmol) were added, and stirred at RT for 2 hr. Most of the precipitate was removed by filtration, the filtrate was washed with water, dried over Na₂SO₄, and concentrated. Purification by flash chromatography (silica gel, 110 g ISCO, 30-50% EtOAc in hexanes) gave the title compound 3B (102 mg). MS (ES) *m/z* 489 [M+H]⁺.

2-Chloro-4-((1*R*,7*S*,7*aS*)-1,7-dihydroxy-3-oxo-1-(trifluoromethyl)-hexahydro-1*H*-pyrrolizin-2-yl)-3-methylbenzonitrile




[00139] To a plastic vial containing compound 3B (20 mg, 0.041 mmol) in 0.2 mL of THF was added 100 µL of HF pyridine solution (Aldrich), and stirred at RT overnight. It was concentrated and purified on prepHPLC to give the title compound (10.2 mg). MS (ES) *m/z* 375 [M+H]⁺.

[00140] While it is apparent that the embodiments of the invention herein disclosed are well suited to fulfill the objectives stated above, it will be appreciated that numerous modifications and other embodiments may be implemented by those skilled in the art, and it is intended that the appended claims cover all such modifications and embodiments that fall within the true spirit and scope of the present invention.

[00141] A number of references have been cited and the entire disclosures of which are incorporated herein by reference.

WHAT IS CLAIMED IS:

1. A compound according to formula I:



5

I

wherein

R_1 is independently selected from the group consisting of H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, arylalkyl, substituted arylalkyl, CO_2R_4 , $CONR_4R_4$, and CH_2OR_4 ;

R_2 is independently selected from the group consisting of H, alkyl, substituted alkyl, OR_3 , SR_3 , halo, NHR_3 , $NHCOR_4$, $NHCO_2R_4$, $NHCONR_4R_4$ and $NHSO_2R_4$;

R_2' is independently selected from the group consisting of OR_3 , SR_3 , halo, NHR_3 , $NHCOR_4$, $NHCO_2R_4$, $NHCONR_4R_4$ and $NHSO_2R_4$;

R_3 is independently selected from the group consisting of H, alkyl, substituted alkyl, CHF_2 , CF_3 and COR_4 ;

R_4 is independently selected from the group consisting of H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

R_5 is independently selected from the group consisting of H, OR_3 , CF_3 , alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl, wherein R_5 and R_5 taken together can form a double bond with O, S, NR_7 or CR_7R_7 when \parallel is a single bond, it being understood that only one R_5 is present when \parallel is a double bond;

W is CR_6R_6 , SO, or SO_2 ;

R_6 is independently selected from the group consisting of H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl, wherein at least one R_6 is H, or R_6 and R_6 taken together can
5 form a double bond with O, S, NR_7 or CR_7R_7 ;

R_7 is independently selected from the group consisting of H, OR_4 , alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

10 G is selected from the group consisting of aryl, heterocyclo and heteroaryl, wherein said aryl, heterocyclo or heteroaryl is mono- or polycyclic, and is optionally substituted with one or more substituents selected from the group consisting of hydrogen, halo, CN, CF_3 , OR_4 , CO_2R_4 , NR_4R_4 , $CONR_4R_4$, CH_2OR_4 , alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted
15 cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl; and

n is 1 or 2; wherein

the definition of formula I above includes all prodrug esters, stereoisomers and pharmaceutically acceptable salts of formula I.

20

2. The compound according to Claim 1, wherein

R_1 is H or alkyl;

R_2' is OH;

R_5 is alkyl or substituted alkyl;

25 \parallel is a double bond;


W is $C=O$; and

n is 1.

30

3. The compound according to Claim 2, wherein

G is selected from:



wherein

R₈, R₉, R₁₀ and R₁₁ are independently selected from the group consisting of H,
 5 NO₂, CN, CF₃, OR₄, CO₂R₄, NR₄R₄, CONR₄R₄, CH₂OR₄, alkyl, substituted alkyl,
 alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted
 cycloalkyl, arylalkyl, substituted arylalkyl, aryl, substituted aryl, heteroaryl and
 substituted heteroaryl;

A, B, C, D, E and F are independently selected from the group consisting of N
 10 and CR₁₂;

J, K, L, P and Q are independently selected from the group consisting of NR₁₃,
 O, S, SO, SO₂ and CR₁₃R₁₃;

R₁₂ is independently selected from the group consisting of a direct bond and
 R₁₃;

15 R₁₃ is independently selected from the group consisting of H, alkyl, substituted
 alkyl, alkenyl, substituted alkenyl, arylalkyl, substituted arylalkyl, CO₂R₄, CONR₄R₄
 and CH₂OR₄; and

z is 0 or 1.

20 4. The compound according to Claim 3, wherein
 R₈ is CN.

5. A pharmaceutical composition, comprising:
 at least one compound according to formula I; and
 25 at least one pharmaceutically acceptable diluent or carrier.

6. A method for preventing, inhibiting or treating the progression or onset
 of diseases or disorders associated with nuclear hormone receptors, comprising:

administering to a patient in need a theapeutically effective amount of a compound according to formula I.

7. The method according to Claim 6, wherein the nuclear hormone
5 receptor is the androgen receptor.

8. The method according to Claim 7, wherein the disease or disorder associated with the androgen receptor is selected from the group consisting of maintenance of muscle strength and function; reversal or prevention of frailty or age-
10 related functional decline ("ARFD") in the elderly (e.g., sarcopenia); prevention of catabolic side effects of glucocorticoids; prevention and treatment of reduced bone density or growth (e.g., osteoporosis and osteopenia); treatment of chronic fatigue syndrome (CFS); chronic myalgia; treatment of acute fatigue syndrome and muscle loss following elective surgery (e.g., post-surgical rehabilitation); treatment of urinary
15 incontinence, acceleration of wound healing; accelerating bone fracture repair (such as accelerating the recovery of hip fracture patients); treatment of wasting secondary to fractures and wasting in connection with chronic obstructive pulmonary disease (COPD), chronic liver disease, AIDS, weightlessness, cancer cachexia, burn and trauma recovery, chronic catabolic state (e.g., coma), eating disorders (e.g., anorexia)
20 and chemotherapy.

9. The method according to claim 7, further comprising:
a therapeutically effective amount of at least one other therapeutic agent.